

RADIOLABELLED QUINOLINE AND QUINOLINONE DERIVATIVES AND
THEIR USE AS METABOTROPIC GLUTAMATE RECEPTOR LIGANDS.

5 The present invention is concerned with radiolabelled quinoline and quinolinone derivatives showing metabotropic glutamate receptor antagonistic activity, in particular mGlu1 receptor activity, and their preparation ; it further relates to compositions comprising them, as well as their use in a diagnostic method, in particular for marking and identifying metabotropic glutamate receptor sites and for imaging an organ.

10

Introduction

The neurotransmitter glutamate is considered to be the major excitatory neurotransmitter in the mammalian central nervous system. The binding of this neurotransmitter to metabotropic glutamate receptors (mGluRs), which are a subfamily
15 of the G-protein-coupled receptors and which comprise 8 distinct subtypes of mGluRs, namely mGluR1 through mGluR8, activates a variety of intracellular second messenger systems. The mGluRs can be divided into 3 groups based on amino acid sequence homology, the second messenger system utilized by the receptors and the pharmacological characteristics. Group I mGluRs, which comprises mGluR subtype 1
20 and 5, couple to phospholipase C and their activation leads to intracellular calcium-ion mobilization. Group II mGluRs (mGluR2 and 3) and group III mGluRs (mGluR4, 6, 7 and 8) couple to adenyl cyclase and their activation causes a reduction in second messenger cAMP and as such a dampening of the neuronal activity. Treatment with Group I mGluR antagonists has been shown to translate in the parasynapsis into a
25 reduced release of neurotransmitter glutamate and to decrease the glutamate-mediated neuronal excitation via postsynaptic mechanisms. Since a variety of pathophysiologic processes and disease states affecting the central nervous system are thought to be due to excessive glutamate induced excitation of the central nervous system neurons, Group I mGluR antagonists, in particular mGluR1 antagonists could be therapeutically
30 beneficial in the treatment of central nervous system diseases, in particular in psychiatric and neurological diseases.

However, up to now, no specific mGluR1-ligands were available, a lack severely hampering the study of the mGlu1 receptors, in particular the radioautographic
35 investigations of the unequivocal distribution and abundance of these receptors in brain sections.. For group I, only [³H]glutamate was available so far, being used on rat (Thomsen et al., Brain Res. 619:22-28, 1993) or human (Kingston et al.,

Neuropharmacology 37:277-287, 1998) mGlu1a receptors. For the mGlu1a receptor and the mGlu5 receptor [³H]quisqualate is available, however, said receptor is not specific for the mGlu1 receptor (it also binds to the AMPA receptor) and it is competitive, i.e. it displaces glutamate (Mutel et al., J. Neurochem. 75:2590-2601, 2000).

It has been the goal of this invention to provide suitable specific, in particular non-competitive mGlu1 receptor ligands.

10 The inventors have now found a particular group of compounds that - in a radiolabelled form - provides for suitable specific, in particular non-competitive mGlu1 receptor ligands as well as a method for marking and identifying metabotropic glutamate receptor sites and for imaging an organ.

15 In the framework of this application, the term "specific" means that the ligand binds preferentially to the mGlu1 receptor site. The term "non-competitive" means that the ligand does not or only marginally displaces glutamate bonded to the mGlu1 receptor site.

20 WO 02/28837 discloses the non-radioactive compounds according to the present invention.


WO 99/26927 discloses antagonists of Group I mGluRs for treating neurological diseases and disorders, based – among others – on a quinoline structure.

25 WO 99/03822 discloses bicyclic metabotropic glutamate receptor ligands, none of them based on a quinoline or quinolinone structure.


30 WO 94/27605 discloses 1,2,3,4-tetrahydroquinoline-2,3,4-trione-3 or 4-oximes and use thereof for treating and preventing neuronal loss, neurodegenerative diseases, adverse consequences of the hyperactivity of the excitatory amino acids and anxiety, as well as radiolabelled compounds thereof.

Detailed Description of the invention

35 The present invention concerns the radiolabelled compounds of Formula (I-A)* or (I-B)*



(I-A)*




(I-B)*

an *N*-oxide form, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemical isomeric form thereof, wherein

X represents O; C(R⁶)₂ with R⁶ being hydrogen, aryl or C₁₋₆alkyl optionally substituted with amino or mono- or di(C₁₋₆alkyl)amino; S or N-R⁷ with R⁷ being amino or hydroxy;

R¹ represents C₁₋₆alkyl; aryl; thienyl; quinoliny; cycloC₃₋₁₂alkyl or (cycloC₃₋₁₂alkyl)C₁₋₆alkyl, wherein the cycloC₃₋₁₂alkyl moiety optionally may contain a double bond and wherein one carbon atom in the cycloC₃₋₁₂alkyl moiety may be replaced by an oxygen atom or an NR⁸-moiety with R⁸ being hydrogen, benzyl or C₁₋₆alkyloxycarbonyl; wherein one or more hydrogen atoms in a C₁₋₆alkyl-moiety or in a cycloC₃₋₁₂alkyl-moiety optionally may be replaced by C₁₋₆alkyl, hydroxyC₁₋₆alkyl, haloC₁₋₆alkyl, aminoC₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, arylC₁₋₆alkyloxy, halo, C₁₋₆alkyloxycarbonyl, aryl, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyloxycarbonylamino, halo, piperazinyl, pyridinyl, morpholinyl, thienyl or a bivalent radical of formula -O-, -O-CH₂-O or -O-CH₂-CH₂-O-; or a radical of formula (a-1)



a-1


wherein Z₁ is a single covalent bond, O, NH or CH₂;

Z₂ is a single covalent bond, O, NH or CH₂;

n is an integer of 0, 1, 2 or 3;

and wherein each hydrogen atom in the phenyl ring independently may optionally be replaced by halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy or hydroxyC₁₋₆alkyl;

or X and R¹ may be taken together with the carbon atom to which X and R¹ are attached to form a radical of formula (b-1), (b-2) or (b-3);



R^2 represents hydrogen; halo; cyano; C_{1-6} alkyl; C_{1-6} alkyloxy; C_{1-6} alkylthio;
 C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl; C_{1-6} alkylcarbonyloxy C_{1-6} alkyl;
 C_{2-6} alkenyl; hydroxy C_{2-6} alkenyl; C_{2-6} alkynyl; hydroxy C_{2-6} alkynyl;
tri(C_{1-6} alkyl)silane C_{2-6} alkynyl; amino; mono- or di(C_{1-6} alkyl)amino; mono- or
di(C_{1-6} alkyloxy C_{1-6} alkyl)amino; mono- or di(C_{1-6} alkylthio C_{1-6} alkyl)amino; aryl;
aryl C_{1-6} alkyl; aryl C_{2-6} alkynyl; C_{1-6} alkyloxy C_{1-6} alkylamino C_{1-6} alkyl;
aminocarbonyl optionally substituted with C_{1-6} alkyl, C_{1-6} alkyloxy C_{1-6} alkyl,
 C_{1-6} alkyloxycarbonyl C_{1-6} alkyl or pyridinyl C_{1-6} alkyl;
a heterocycle selected from thienyl, furanyl, pyrrolyl, thiazolyl, oxazolyl,
imidazolyl, isothiazolyl, isoxazolyl, pyrazolyl, pyridyl, pyrazinyl, pyridazinyl,
pyrimidinyl, piperidinyl and piperazinyl, optionally N-substituted with
 C_{1-6} alkyloxy C_{1-6} alkyl, morpholinyl, thiomorpholinyl, dioxanyl or dithianyl ;
a radical $-NH-C(=O)R^9$ wherein R^9 represents
 C_{1-6} alkyl optionally substituted with cyclo C_{3-12} alkyl, C_{1-6} alkyloxy,
 C_{1-6} alkyloxycarbonyl, aryl, aryloxy, thienyl, pyridinyl, mono- or
di(C_{1-6} alkyl)amino, C_{1-6} alkylthio, benzylthio, pyridinylthio or
pyrimidinylthio;
cyclo C_{3-12} alkyl; cyclohexenyl; amino; arylcyclo C_{3-12} alkylamino;
mono-or-di(C_{1-6} alkyl)amino; mono- or
di(C_{1-6} alkyloxycarbonyl C_{1-6} alkyl)amino; mono- or
di(C_{1-6} alkyloxycarbonyl)amino; mono-or di(C_{2-6} alkenyl)amino; mono- or
di(aryl C_{1-6} alkyl)amino; mono- or diarylamino; aryl C_{2-6} alkenyl;
furanyl C_{2-6} alkenyl; piperidinyl; piperazinyl; indolyl; furyl; benzofuryl;
tetrahydrofuryl; indenyl; adamantyl; pyridinyl; pyrazinyl; aryl;
aryl C_{1-6} alkylthio or a radical of formula (a-1) ;
a sulfonamid $-NH-SO_2-R^{10}$ wherein R^{10} represents C_{1-6} alkyl, mono- or poly
halo C_{1-6} alkyl, aryl C_{1-6} alkyl, aryl C_{2-6} alkenyl, aryl, quinolinyl, isoxazolyl
or di(C_{1-6} alkyl)amino;
 R^3 and R^4 each independently represent hydrogen; halo; hydroxy; cyano; C_{1-6} alkyl;
 C_{1-6} alkyloxy; C_{1-6} alkyloxy C_{1-6} alkyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl;

C₂₋₆alkenyl; hydroxyC₂₋₆alkenyl; C₂₋₆alkynyl; hydroxyC₂₋₆alkynyl;
tri(C₁₋₆alkyl)silaneC₂₋₆alkynyl; amino; mono- or di(C₁₋₆alkyl)amino; mono- or
di(C₁₋₆alkyloxyC₁₋₆alkyl)amino; mono- or di(C₁₋₆alkylthioC₁₋₆alkyl)amino; aryl;
morpholinylC₁₋₆alkyl or piperidinylC₁₋₆alkyl ; or

5 R² and R³ may be taken together to form -R²-R³-, which represents a bivalent radical of
formula -(CH₂)₃-, -(CH₂)₄-, -(CH₂)₅-, -(CH₂)₆-, -CH=CH-CH=CH-,
-Z₄-CH=CH-, -CH=CH-Z₄-, -Z₄-CH₂-CH₂-CH₂-, -CH₂-Z₄-CH₂-CH₂-,
-CH₂-CH₂-Z₄-CH₂-,
-CH₂-CH₂-CH₂-Z₄-, -Z₄-CH₂-CH₂-, -CH₂-Z₄-CH₂- or -CH₂-CH₂-Z₄-, with Z₄ being
10 O, S, SO₂ or NR¹¹ wherein R¹¹ is hydrogen, C₁₋₆alkyl, benzyl or
C₁₋₆alkyloxycarbonyl; and wherein each bivalent radical is optionally substituted
with C₁₋₆alkyl.

or R³ and R⁴ may be taken together to form a bivalent radical of formula
-CH=CH-CH=CH- or -CH₂-CH₂-CH₂-CH₂- ;

15 R⁵ represents hydrogen; cycloC₃₋₁₂alkyl; piperidinyl; oxo-thienyl; tetrahydrothienyl,
arylC₁₋₆alkyl; C₁₋₆alkyloxyC₁₋₆alkyl; C₁₋₆alkyloxycarbonylC₁₋₆alkyl or C₁₋₆alkyl
optionally substituted with a radical C(=O)NR_xR_y, in which R_x and R_y, each
independently are hydrogen, cycloC₃₋₁₂alkyl, C₂₋₆alkynyl or C₁₋₆alkyl optionally
substituted with cyano, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, furanyl, pyrrolidinyl,
20 benzylthio, pyridinyl, pyrrolyl or thienyl;

Y represents O or S;

or Y and R⁵ may be taken together to form =Y-R⁵- which represents a radical of
formula

-CH=N-N= (c-1);
25 -N=N-N= (c-2); or
-N-CH=CH- (c-3);

aryl represents phenyl or naphthyl optionally substituted with one or more substituents
selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, phenyloxy, nitro, amino, thio,
C₁₋₆alkylthio, haloC₁₋₆alkyl, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy,
30 hydroxyC₁₋₆alkyl, C₁₋₆alkyloxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono-or
di(C₁₋₆alkyl)amino; mono-or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, cyano, -CO-R¹²,
-CO-OR¹³, -NR¹³SO₂R¹², -SO₂-NR¹³R¹⁴, -NR¹³C(O)R¹², -C(O)NR¹³R¹⁴, -SOR¹²,
-SO₂R¹²; wherein each R¹², R¹³ and R¹⁴ independently represent C₁₋₆alkyl;
cycloC₃₋₆alkyl; phenyl; phenyl substituted with halo, hydroxy, C₁₋₆alkyl,
35 C₁₋₆alkyloxy, haloC₁₋₆alkyl, polyhaloC₁₋₆alkyl, furanyl, thienyl, pyrrolyl,
imidazolyl, thiazolyl or oxazolyl;

and when the $R^1-C(=X)$ moiety is linked to another position than the 7 or 8 position, then said 7 and 8 position may be substituted with R^{15} and R^{16} wherein either one or both of R^{15} and R^{16} represents C_{1-6} alkyl, C_{1-6} alkyloxy or R^{15} and R^{16} taken together may form a bivalent radical of formula $-CH=CH-CH=CH-$.

5

As used in the foregoing definitions and hereinafter C_{1-6} alkyl as a group or part of a group encompasses the straight and branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as, for example, methyl, ethyl, propyl, butyl, pentyl or hexyl; C_{2-6} alkenyl as a group or part of a group encompasses the straight and branched chain hydrocarbon radicals having from 2 to 6 carbon atoms and having a double bond such as ethenyl, propenyl, butenyl, pentenyl, hexenyl, 3-methylbutenyl and the like; C_{2-6} alkynyl as a group or part of a group defines straight or branched chain hydrocarbon radicals having from 2 to 6 carbon atoms and having a triple bond such as ethynyl, propynyl, butynyl, pentynyl, hexynyl, 3-methylbutynyl and the like;

10

15

cyclo C_{3-6} alkyl encompasses monocyclic alkyl ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl; cyclo C_{3-12} alkyl encompasses mono-, bi- or tricyclic alkyl ring structures and is generic to for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, norbornanyl, adamantyl.

20

The term halo is generic to fluoro, chloro, bromo and iodo. As used in the foregoing and hereinafter, polyhalo C_{1-6} alkyl as a group or part of a group is defined as mono- or polyhalosubstituted C_{1-6} alkyl, in particular methyl with one or more fluoro atoms, for example, difluoromethyl or trifluoromethyl. In case more than one halogen atoms are attached to an alkyl group within the definition of polyhalo C_{1-6} alkyl, they may be the same or different.

25

When any variable, e.g. aryl, occurs more than one time in any constituent, each definition is independent.

30

When any bond is drawn into a ring structure, it means that the corresponding substituent may be linked to any atom of said ring structure. This means for instance that the $R^1-C(=X)$ moiety may be linked to the quinoline or quinolinone moiety in position 5, 6, 7, 8 but also position 3 or position 4.

35

By the term "radiolabelled compound" is meant any compound according to Formula (I-A)* or (I-B)*, an *N*-oxide form, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, which contains at least

one radioactive atom. In the framework of this application, compounds which do not contain a radio-active atom are denoted without an asterisk to their formula number, compounds which contain a radio-active atom are denoted with an asterisk to their formula number. Compounds can be labelled with either positron or gamma emitting radionuclides. For radioligand-binding techniques (membrane receptor assay), the $[^3\text{H}]$ -atom or the $[^{125}\text{I}]$ -atom is the atom of choice. For imaging, the most commonly used positron emitting (PET) radionuclides are ^{11}C , ^{18}F , ^{15}O and ^{13}N , all of which are accelerator produced and have half-lives of 20, 100, 2 and 10 minutes respectively. Since the half-lives of these radionuclides are so short, it is only feasible to use them at institutions which have an accelerator on site for their production, thus limiting their use. The most widely used of these are ^{18}F , $^{99\text{m}}\text{Tc}$, ^{201}Tl and ^{123}I .

In particular, the radioactive atom is selected from the group of hydrogen, carbon, nitrogen, sulfur, oxygen and halogen. Preferably, the radioactive atom is selected from the group of hydrogen, carbon and halogen.

In particular, the radioactive atom is selected from the group of ^3H , ^{11}C , ^{18}F , ^{122}I , ^{123}I , ^{125}I , ^{131}I , ^{75}Br , ^{76}Br , ^{77}Br and ^{82}Br . Preferably, the radioactive atom is selected from the group of ^3H , ^{11}C and ^{18}F .

By the term "compound according to the invention" is meant a compound according to Formula (I-A)* or (I-B)*, an *N*-oxide form, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof.

For *in vivo* use, salts of the compounds of Formula (I-A)* and (I-B)* are those wherein the counter ion is pharmaceutically acceptable. However, salts of acids and bases which are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention. With the term "in vivo" is meant any use of the compounds according to the invention whereby said compounds are administered to live animals.

The pharmaceutically acceptable addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid addition salt forms which the compounds of Formula (I-A)* and (I-B)* are able to form. The latter can conveniently be obtained by treating the base form with such appropriate acids as inorganic acids, for example, hydrohalic acids, e.g. hydrochloric, hydrobromic and the like; sulfuric acid;

nitric acid; phosphoric acid and the like; or organic acids, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 2-oxopropanoic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids. Conversely the salt form can be converted by treatment with alkali into the free base form.

The compounds of Formula (I-A)* and (I-B)* containing acidic protons may be converted into their therapeutically active non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. primary, secondary and tertiary aliphatic and aromatic amines such as methylamine, ethylamine, propylamine, isopropylamine, the four butylamine isomers, dimethylamine, diethylamine, diethanolamine, dipropylamine, diisopropylamine, di-n-butylamine, pyrrolidine, piperidine, morpholine, trimethylamine, triethylamine, tripropylamine, quinuclidine, pyridine, quinoline and isoquinoline, the benzathine, *N*-methyl-D-glucamine, 2-amino-2-(hydroxymethyl)-1,3-propanediol, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like. Conversely the salt form can be converted by treatment with acid into the free acid form.

The term "addition salt" also comprises the hydrates and solvent addition forms which the compounds of Formula (I-A)* and (I-B)* are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of Formula (I-A)* and (I-B)* are able to form by reaction between a basic nitrogen of a compound of Formula (I-A)* or (I-B)* and an appropriate quaternizing agent, such as, for example, an optionally substituted alkylhalide, arylhalide or arylalkylhalide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl *p*-toluenesulfonates. A quaternary amine has a positively charged nitrogen. Pharmaceutically acceptable counter ions include chloro, bromo, iodo, trifluoroacetate and acetate. The counter ion of choice can be introduced using ion exchange resins.

It will be appreciated that some of the compounds according to the invention may contain one or more centers of chirality and exist as stereochemically isomeric forms.

5 The term "stereochemically isomeric forms" as used hereinbefore defines all the possible stereoisomeric forms which the compounds according to the invention or physiologically functional derivatives may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereoisomeric forms, said mixtures containing all diastereomers and enantiomers of the
10 basic molecular structure as well as each of the individual isomeric forms of the compounds according to the invention, substantially free, i.e. associated with less than 10 %, preferably less than 5 %, in particular less than 2% and most preferably less than 1 % of the other isomers. Stereochemically isomeric forms of the compounds according to the invention are obviously intended to be embraced within the scope of
15 the present invention. The same applies to the intermediates as described herein, used to prepare end products of the compounds according to the invention.

The terms *cis* and *trans* are used herein in accordance with Chemical Abstracts nomenclature.

20 In some compounds according to the invention and in the intermediates used in their preparation, the absolute stereochemical configuration has not been determined. In these cases, the stereoisomeric form which was first isolated is designated as "A" and the second as "B", without further reference to the actual stereochemical configuration.
25 However, said "A" and "B" stereoisomeric forms can be unambiguously characterized by physicochemical characteristics such as their optical rotation in case "A" and "B" have an enantiomeric relationship. A person skilled in the art is able to determine the absolute configuration of such compounds using art-known methods such as, for example, X-ray diffraction. In case "A" and "B" are stereoisomeric mixtures, they can
30 be further separated whereby the respective first fractions isolated are designated "A1" and "B1" and the second as "A2" and "B2", without further reference to the actual stereochemical configuration.

The *N*-oxide forms of the present compounds are meant to comprise the compounds of
35 formula (I-A)* and (I-B)* wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide.


Some of the compounds according to the invention may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention. Of special interest are those compounds of formula (I-A)* and (I-B)* which are stereochemically pure.

5

An interesting group of compounds are those compounds of formula (I-A)* and (I-B)* wherein

X represents O; C(R⁶)₂ with R⁶ being hydrogen or aryl ; or N-R⁷ with R⁷ being amino or hydroxy;

10 R¹ represents C₁₋₆alkyl, aryl; thienyl; quinoliny; cycloC₃₋₁₂alkyl or (cycloC₃₋₁₂alkyl)C₁₋₆alkyl, wherein the cycloC₃₋₁₂alkyl moiety optionally may contain a double bond and wherein one carbon atom in the cycloC₃₋₁₂alkyl moiety may be replaced by an oxygen atom or an NR⁸-moiety with R⁸ being benzyl or C₁₋₆alkyloxycarbonyl ; wherein one or more hydrogen atoms in a C₁₋₆alkyl-moiety
15 or in a cycloC₃₋₁₂alkyl-moiety optionally may be replaced by C₁₋₆alkyl, haloC₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, arylC₁₋₆alkyloxy, halo, aryl, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyloxycarbonylamino, halo, piperazinyl, pyridinyl, morpholinyl, thienyl or a bivalent radical of formula -O- or -O-CH₂-CH₂-O-; or a radical of formula (a-1)



a-1

20

wherein Z₁ is a single covalent bond, O or CH₂;


Z₂ is a single covalent bond, O or CH₂;

n is an integer of 0, 1, or 2 ;


25

and wherein each hydrogen atom in the phenyl ring independently may optionally be replaced by halo or hydroxy;


or X and R¹ may be taken together with the carbon atom to which X and R¹ are attached to form a radical of formula (b-1), (b-2) or (b-3);



b-1



b-2



b-3

30

R^2 represents hydrogen; halo; cyano; C_{1-6} alkyl; C_{1-6} alkyloxy; C_{1-6} alkylthio;
 C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl; C_{2-6} alkenyl; hydroxy C_{2-6} alkenyl;
 C_{2-6} alkynyl; hydroxy C_{2-6} alkynyl; tri(C_{1-6} alkyl)silane C_{2-6} alkynyl; amino; mono- or
5 di(C_{1-6} alkyl)amino; mono- or di(C_{1-6} alkyloxy C_{1-6} alkyl)amino; mono- or
di(C_{1-6} alkylthio C_{1-6} alkyl)amino; aryl; aryl C_{1-6} alkyl; aryl C_{2-6} alkynyl;
 C_{1-6} alkyloxy C_{1-6} alkylamino C_{1-6} alkyl;
aminocarbonyl optionally substituted with C_{1-6} alkyloxycarbonyl C_{1-6} alkyl ;
a heterocycle selected from thienyl, furanyl, thiazolyl and piperidinyl, optionally
10 N-substituted with morpholinyl or thiomorpholinyl;
a radical $-NH-C(=O)R^9$ wherein R^9 represents C_{1-6} alkyl optionally substituted with
cyclo C_{3-12} alkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, aryl, aryloxy, thienyl,
pyridinyl, mono- or di(C_{1-6} alkyl)amino, C_{1-6} alkylthio, benzylthio, pyridinylthio or
pyrimidinylthio; cyclo C_{3-12} alkyl; cyclohexenyl; amino; arylcyclo C_{3-12} alkylamino;
15 mono-or-di(C_{1-6} alkyl)amino; mono- or di(C_{1-6} alkyloxycarbonyl C_{1-6} alkyl)amino;
mono- or di(C_{1-6} alkyloxycarbonyl)amino; mono-or di(C_{2-6} alkenyl)amino; mono- or
di(aryl C_{1-6} alkyl)amino; mono- or diarylamino; aryl C_{2-6} alkenyl; furanyl C_{2-6} alkenyl;
piperidinyl; piperazinyl; indolyl; furyl; benzofuryl; tetrahydrofuryl; indenyl;
adamantyl; pyridinyl; pyrazinyl; aryl or a radical of formula (a-1) ;
20 a sulfonamid $-NH-SO_2-R^{10}$ wherein R^{10} represents C_{1-6} alkyl, mono- or poly
halo C_{1-6} alkyl, aryl C_{1-6} alkyl or aryl;
 R^3 and R^4 each independently represent hydrogen; C_{1-6} alkyl; C_{1-6} alkyloxy C_{1-6} alkyl;
 C_{1-6} alkyloxycarbonyl; or
 R^2 and R^3 may be taken together to form $-R^2-R^3-$, which represents a bivalent radical of
25 formula $-(CH_2)_4-$, $-(CH_2)_5-$, $-Z_4-CH=CH-$, $-Z_4-CH_2-CH_2-CH_2-$ or $-Z_4-CH_2-CH_2-$,
with Z_4 being O, S, SO_2 or NR^{11} wherein R^{11} is hydrogen, C_{1-6} alkyl, benzyl or
 C_{1-6} alkyloxycarbonyl; and wherein each bivalent radical is optionally substituted
with C_{1-6} alkyl;
or R^3 and R^4 may be taken together to form a bivalent radical of formula
30 $-CH=CH-CH=CH-$ or $-CH_2-CH_2-CH_2-CH_2-$;
 R^5 represents hydrogen; piperidinyl; oxo-thienyl; tetrahydrothienyl, aryl C_{1-6} alkyl;
 C_{1-6} alkyloxycarbonyl C_{1-6} alkyl or C_{1-6} alkyl optionally substituted with a radical
 $C(=O)NR_xR_y$, in which R_x and R_y , each independently are hydrogen,
cyclo C_{3-12} alkyl, C_{2-6} alkynyl or C_{1-6} alkyl optionally substituted with cyano,
35 C_{1-6} alkyloxy or C_{1-6} alkyloxycarbonyl;
Y represents O or S;

or Y and R⁵ may be taken together to form =Y-R⁵- which represents a radical of formula

-CH=N-N= (c-1); or

-N=N-N= (c-2);

5 aryl represents phenyl or naphthyl optionally substituted with one or more substituents selected from halo, C₁₋₆alkyloxy, phenyloxy, mono-or di(C₁₋₆alkyl)amino and cyano;

and when the R¹-C(=X) moiety is linked to another position than the 7 or 8 position, then said 7 and 8 position may be substituted with R¹⁵ and R¹⁶ wherein either one or
10 both of R¹⁵ and R¹⁶ represents C₁₋₆alkyl or R¹⁵ and R¹⁶ taken together may form a bivalent radical of formula -CH=CH-CH=CH-.

A further most interesting group of compounds comprises those compounds of formula (I-A)* and (I-B)* wherein X represents O;

15 R¹ represents C₁₋₆alkyl; cycloC₃₋₁₂alkyl or (cycloC₃₋₁₂alkyl)C₁₋₆alkyl, wherein one or more hydrogen atoms in a C₁₋₆alkyl-moiety or in a cycloC₃₋₁₂alkyl-moiety optionally may be replaced by C₁₋₆alkyloxy, aryl, halo or thienyl;

R² represents hydrogen; halo; C₁₋₆alkyl or amino;

R³ and R⁴ each independently represent hydrogen or C₁₋₆alkyl; or

20 R² and R³ may be taken together to form -R²-R³-, which represents a bivalent radical of formula -Z₄-CH₂-CH₂-CH₂- or -Z₄-CH₂-CH₂- with Z₄ being O or NR¹¹ wherein R¹¹ is C₁₋₆alkyl; and wherein each bivalent radical is optionally substituted with C₁₋₆alkyl;

or R³ and R⁴ may be taken together to form a bivalent radical of formula
25 -CH₂.CH₂-CH₂-CH₂- ;

R⁵ represents hydrogen;


Y represents O; and

aryl represents phenyl optionally substituted with halo.

30 A further interesting group of compounds comprises those compounds of formula (I-A)* and (I-B)* wherein the R¹-C(=X) moiety is linked to the quinoline or quinolinone moiety in position 6.

Especially interesting are the following radioactive compounds :

35



All compounds according to the invention show a moderate to strong mGluR1 activity. Such activity is among others attributed to the specific binding of said compound to the mGlu1 receptor, which makes the compounds useful in a diagnostic method, e.g. for labeling and detecting mGlu1 receptor sites. The invention therefore also relates to a radiolabelled compound according to the invention for use in a diagnostic method.

First preferred embodiment (gamma emitting radionuclide)

According to a first preferred embodiment, the radiolabelled compound comprises at least one ^3H -atom or one ^{125}I -atom. A ^3H -atom is conveniently introduced by partially or completely substituting one or more non-radioactive ^1H -hydrogen atoms in the molecule by their radioactive isotopes. The choice of whether a ^3H or ^{125}I radioligand will be used may depend in part on the availability of liquid scintillation counters (LSC), which are fairly expensive. ^{125}I ligands can be quantified using either a γ -counter or an LSC, whereas ^3H ligands necessitate the use of an LSC.

The radiolabelled compound comprising at least one [^3H]-atom or one [^{125}I]-atom is advantageously used in radioligand-binding techniques, in particular in *in vitro* membrane receptor assays for marking or identifying a mGlu1 receptor in biological material.

The radiolabelled compounds comprising at least one [^3H]-atom or one [^{125}I]-atom is also advantageously used in *in vivo* mGlu1 receptor autoradiography of the brain since the compounds according to the invention have the advantageous and unexpected ability to readily cross the blood-brain barrier.

The invention therefore also relates to a radiolabelled compound according to the invention used in a diagnostic method which consists of marking or identifying a mGlu1 receptor in biological material, as well as the use of the compounds according to the invention for the manufacture of a diagnostic tool for marking or identifying an mGlu1 receptor in biological material, whether *in vivo* or *in vitro*.

In the framework of this application, by the term "biological material" is meant to include any material which has a biological origin. In particular, this relates to tissue samples, plasma fluids, body fluids, body parts and organs originating from warm-blooded animals and warm-blooded animals *per se*, in particular humans.

Basic experiments that are performed using the membrane assay system for marking or identifying a mGlu1 receptor in biological material are : saturation experiments, inhibition experiments, association kinetic experiments and dissociation kinetic experiments. These methods are applicable to most neurotransmitter and hormone receptor systems, including the mGluR1-system (Methods for Neurotransmitter Receptor Analysis, ed. by Henry I. Yamamura et al., Raven Press Ltd., New York, 1990). To this end, the radiolabelled compound is administered to the biological material to mark the mGlu1 receptors and the emissions from the radiolabelled compound are detected to identify the amount or location of the mGlu1 receptors, for instance for *ex vivo* receptor autoradiography.

The radiolabelled compounds according to the invention comprising at least one [^3H]-atom or one [^{125}I]-atom are also useful as agents for screening whether a test compound has the ability to occupy or bind to a mGlu1 receptor site. The degree to which a test compound will displace a compound according to the invention from the mGlu1 receptor site will show the test compound ability to occupy or bind to a mGlu1

receptor and therefore act as either an agonist, an antagonist or a mixed agonist/antagonist of a mGlu1 receptor.


The radiolabelled compounds according to the invention comprising at least one $[^3\text{H}]$ -atom or one $[^{125}\text{I}]$ -atom are advantageously prepared by substituting a halogen atom with a tritium atom, as is documented in the Experimental Section below.

Second Preferred Embodiment (positron emitting radionuclide)

In a second preferred embodiment, the radiolabelled compound comprises at least one radioactive carbon or halogen atom. In principle, any compound according to Formula (I) containing a carbon or halogen atom is prone for radiolabel ling by replacing the carbon or halogen atom by a suitable radioactive isotope or by making the compounds according to Formula (I) using radioactively-labelled reagentia. Suitable halogen radioisotopes to this purpose are radioactive carbon, e.g. $[^{11}\text{C}]$; radioactive iodides, e.g. $[^{122}\text{I}]$, $[^{123}\text{I}]$, $[^{131}\text{I}]$; radioactive bromides, e.g. $[^{75}\text{Br}]$, $[^{76}\text{Br}]$, $[^{77}\text{Br}]$ and $[^{82}\text{Br}]$; and radioactive fluorides, e.g. $[^{18}\text{F}]$. Preferred radiolabelled compounds are those compounds of Formula (I-A)* and (I-B)*, wherein R^1 comprises a radioactive carbon or halo atom, especially $[^{11}\text{C}]$, $[^{18}\text{F}]$, $[^{123}\text{I}]$, $[^{75}\text{Br}]$, $[^{76}\text{Br}]$ or $[^{77}\text{Br}]$.

Preparation of the radioactive compounds


The introduction of a radioactive halogen atom can be performed by a suitable reaction such as depicted below



in which all substituents in Formula (I-A-a) and (I-A-a)* are defined as in Formula (I-A)* and halo is a halogen atom. A suitable compound (I-A-a) is reacted with $\text{H}[^{18}\text{F}]$ such that the halogen atom present on the quinoline ring is displaced by a nucleophilic displacement reaction with the radioactive $[^{18}\text{F}]$ atom.

For obtaining radiolabelled compounds according to Formula (I-B)*, radiolabel ling can be performed on an equivalent way, for instance by way of a reaction scheme as

depicted below. Obviously, also compounds according to Formula (I-A)* can be obtained in an equivalent way, i.e. by way of labelling an R¹ substituent.




5

Other methods for tritium-labelling are disclosed e.g. by Peng et al. Fusion Technology, American Nuclear Society, **21(2)**:307-311, 1992 and by Brundish et al. Journal of Labelled Compounds and Radiopharmaceuticals **25(12)**:1361-1369 (1988).

10 The introduction of a radioactive [¹¹C] can be performed using the reaction scheme below in which a suitable compound (I-A-a) is first stannylated after which the radioactive [¹¹C] is introduced using e.g. a palladium catalyzed "Stille-type" coupling reaction using [¹¹C]methyl iodide (Scott, W.J.; Crisp, G.T.; Stille, J.K. *J. Am. Chem. Soc.*, **1984**, *106*, 4630).

15



In Formula (I-A-a), (I-A-b) and (I-A-c)*, all substituents have the same meaning as defined in Formula (I-A)*, halo is a halogen atom and R¹⁷ is methyl or butyl.

20

Because of their unexpected property to penetrate readily the blood-brain barrier, the radiolabelled compounds comprising a radioactive halogen atom are advantageously administered *in vivo*, in an appropriate composition to an animal, especially a warm-blooded animal, and the location of said radiolabelled compounds is detected using

25 imaging techniques, such as, for instance, Single Photon Emission Computerized Tomography (SPECT) or Positron Emission Tomography (PET) and the like. In this manner the distribution of mGlu1 receptor sites throughout the body can be detected


and organs containing mGlu1 receptor sites such as, for example, the brain, can be visualized by the imaging techniques mentioned hereinabove. This process of imaging an organ by administering a radiolabelled compound of Formula (I-A)* or (I-A)*, which bind to the mGlu1 receptor sites and detecting the emissions from the radioactive compound also constitutes an aspect of the present invention.

The application of the compounds of Formula (I-A)* and (I-B)* in the above described techniques constitutes a further aspect of the present invention. The invention in particular relates to the use of the compounds according to the invention for the manufacture of a diagnostic tool for use in PET. For use in PET, most preferred are radiolabelled compounds according to the invention, in which a ^{18}F is incorporated (US 4,931,270 by Horn et al., published June 5, 1990).


Preparation of the non-radioactive compounds

The non-radioactive compounds according to the invention may be produced in a number of ways.


In order to simplify the structural representation of some of the present compounds and intermediates in the following preparation procedures, the quinoline or the quinolinone moiety will hereinafter be represented by the symbol Q.




The compounds of formula (I-A) or (I-B), wherein X represents O, said compounds being represented by formula (I_{A/B}-a), can be prepared by oxidizing an intermediate of formula (II) in the presence of a suitable oxidizing agent, such as potassium permanganate, and a suitable phase-transfer catalyst, such as tris(dioxa-3,6-heptyl)amine, in a suitable reaction-inert solvent, such as for example dichloromethane.




Compounds of formula (I_{A/B}-a) may also be prepared by reacting an intermediate of formula (III) with an intermediate of formula (IV), wherein W₁ represents a halo atom, e.g. bromo, in the presence of butyl lithium and a suitable reaction-inert solvent, such as for example tetrahydrofuran.




Alternatively, compounds of formula (I_{A/B}-a) may also be prepared by reacting an intermediate of formula (V) with an intermediate of formula (IV) in the presence of butyl lithium and a suitable reaction-inert solvent, such as for example tetrahydrofuran.



Compounds of formula (I_{A/B}-a), wherein the R¹ substituent is linked to the carbonyl moiety via an oxygen atom, said R¹ substituent being represented by O-R^{1a} and said compounds by formula (I_{A/B}-a-1), can be prepared by reacting an intermediate of formula (VI) with an intermediate of formula (VII) in the presence of a suitable acid, such as sulfuric acid.



Compounds of formula (I-A), wherein R² represents methylcarbonyl, said compounds being represented by formula (I-A-1), can be prepared by reacting an intermediate of formula (VIII) in the presence of a suitable acid, such as hydrochloric acid, and a suitable reaction-inert solvent, such as for example tetrahydrofuran.



The compounds of formula (I) may also be converted into each other following art-known transformations.

Compounds of formula (I-A) wherein R^2 is a halo atom, such as chloro, can be converted into a compound of formula (I-A), wherein R^2 is another halo atom, such as fluoro or iodo, by reaction with a suitable halogenating agent, such as for example
5 potassium fluoride or sodium iodide, in the presence of a suitable reaction-inert solvent, e.g. dimethyl sulfoxide or acetonitrile and optionally in the presence of acetyl chloride.

Compounds of formula (I-A), wherein R^2 is a suitable leaving group, such as a halo atom, e.g. chloro, iodo, said leaving group being represented by W^2 and said
10 compounds by (I-A-2), can be converted into a compound of formula (I-A) wherein R^2 is cyano, said compound being represented by formula (I-A-3), by reaction with a suitable cyano-introducing agent, such as for example trimethylsilanecarbonitrile, in the presence of a suitable base such as *N,N*-diethylethanamine and a suitable catalyst, such as for example tetrakis(triphenylphosphine)palladium.

15 Compounds of formula (I-A-2) can also be converted into a compound of formula (I-A-4) by reaction with C_{2-6} alkynyltri(C_{1-6} alkyl)silane in the presence of CuI, an appropriate base, such as for example *N,N*-diethylethanamine, and an appropriate catalyst, such as for example tetrakis(triphenylphosphine)palladium. Compounds of
20 formula (I-A-4) can on their turn be converted into a compound of formula (I-A-5) by reaction with potassium fluoride in the presence of a suitable acid such as acetic acid, or by reaction with a suitable base, such as potassium hydroxide, in the presence of a suitable reaction-inert solvent, such as an alcohol, e.g. methanol and the like.

25 Compounds of formula (I-A-2) can also be converted into a compound of formula (I-A-6) by reaction with an intermediate of formula (IX) in the presence of CuI, a suitable base, such as for example *N,N*-diethylethanamine, and a suitable catalyst such as tetrakis(triphenylphosphine)palladium.


30 Compounds of formula (I-A-2) can also be converted into a compound wherein R^2 is C_{1-6} alkyl, said compound being represented by formula (I-A-8) in the presence of a suitable alkylating agent, such as for example $Sn(C_{1-6}alkyl)_4$, or into a compound wherein R^2 is C_{2-6} alkenyl, said compound being represented by formula (I-A-9) in the presence of a suitable alkenylating agent, such as for example $Sn(C_{2-6}alkenyl)$
35 $(C_{1-6}alkyl)_3$, both reactions in the presence of a suitable catalyst, such as for example tetrakis(triphenylphosphine)palladium and a reaction-inert solvent, such as for example toluene or dioxane.

Compounds of formula (I-A-2) can also be converted into a compound of formula (I-A-7) wherein Z represents O or S, by reaction with an intermediate of formula (X) optionally in the presence of a suitable base such as dipotassium carbonate and a
5 reaction-inert solvent, such as *N,N*-dimethyl formamide.


Compounds of formula (I-A-2) can also be converted into a compound of formula (I-A), wherein R^2 is C_{1-6} alkyloxycarbonyl, said compound being represented by formula (I-A-10) and a compound of formula (I-A), wherein R^2 is hydrogen, said
10 compound being represented by formula (I-A-11), by reaction with a suitable alcohol of formula $C_{1-6}alkylOH$ and CO in the presence of a suitable catalyst, such as for example palladium(II)acetate, triphenylphosphine, a suitable base such as dipotassium carbonate and a reaction-inert solvent, such as *N,N*-dimethylformamide.

15 Compounds of formula (I-A-11) can also be prepared by reacting a compound of formula (I-A-2) with Zn in the presence of a suitable acid such as acetic acid.


Compounds of formula (I-A-2) can also be converted into a compound of formula (I-A), wherein R^2 is aminocarbonyl substituted with C_{1-6} alkyloxycarbonyl $C_{1-6}alkyl$,
20 said compound being represented by formula (I-A-12), by reaction with an intermediate of formula $H_2N-C_{1-6}alkyl-C(=O)-O-C_{1-6}alkyl$ in the presence of CO, a suitable catalyst such as tetrakis(triphenylphosphine)palladium, a suitable base, such as for example *N,N*-diethylethanamine, and a suitable reaction-inert solvent, such as for example
25 toluene.



Compounds of formula (I-A-2) can also be converted into a compound of formula (I-A) wherein R^2 is aryl or a heterocycle selected from the group described in the definition of R^2 hereinabove, said R^2 being represented by R^{2a} and said compound by formula (I-A-13) by reaction with an intermediate of formula (XI), (XII) or (XIII) in the presence of a suitable catalyst such as for example tetrakis(triphenylphosphine)palladium and a suitable reaction-inert solvent, such as for example dioxane.




Compounds of formula (I-A-2) can also be converted into a compound of formula (I-B), wherein Y and R^5 are taken together to form a radical of formula (b-1) or (b-2), said compound being represented by formula (I-B-1) or (I-B-2), by reaction with hydrazinecarboxaldehyde or sodium azide in a suitable reaction-inert solvent, such as an alcohol, e.g. butanol, or *N,N*-dimethylformamide.




Compounds of formula (I-A-11) can be converted into the corresponding *N*-oxide, represented by formula (I-A-14), by reaction with a suitable peroxide, such as 3-chloro-benzenecarboxoperoxoic acid, in a suitable reaction-inert solvent, such as for


example methylene chloride. Said compound of formula (I-A-14) can further be converted into a compound of formula (I-B), wherein R^5 is hydrogen, said compound being represented by formula (I-B-3), by reaction with 4-methyl-benzene sulfonyl chloride in the presence of a suitable base, such as for example dipotassium carbonate and a suitable reaction-inert solvent, such as for example methylene chloride.




Compounds of formula (I-B-3) can also be prepared from a compound of formula (I-A), wherein R^2 is C₁₋₆alkyloxy, said compound being represented by formula (I-A-15), by reaction with a suitable acid, such as hydrochloric acid, in the presence of a suitable reaction-inert solvent, such as for example tetrahydrofuran.



Compounds of formula (I-B-3) can be converted into a compound of formula (I-B), wherein R^5 represents C₁₋₆alkyl, said compound being represented by formula (I-B-4), by reaction with an appropriate alkylating agent, such as for example an intermediate of formula (XIV), wherein W₃ represents a suitable leaving group such as a halo atom e.g. iodo, in the presence of potassium tert. butoxide and in the presence of a suitable reaction-inert solvent, such as for example tetrahydrofuran.




Compounds of formula (I-B-3) can also be converted into a compound of formula (I-B), wherein R⁵ is C₁₋₆alkyloxycarbonylC₁₋₆alkyl or arylC₁₋₆alkyl, said R⁵ being represented by R^{5a} and said compound being represented by formula (I-B-5), by reaction with an intermediate of formula (XV), wherein W₄ represents a suitable leaving group, such as a halo atom, e.g. bromo, chloro and the like, in the presence of a suitable base, such as for example sodium hydride and a suitable reaction-inert solvent, such as for example *N,N*-dimethylformamide.




Compounds of formula (I-A-2) can also be converted into a compound of formula (I-B), wherein R⁵ is hydrogen and Y is S, said compound being represented by formula (I-B-6), by reaction with H₂N-C(=S)-NH₂ in the presence of a suitable base, such as

potassium hydroxide, and a suitable reaction-inert solvent, such as an alcohol, for example ethanol, or water. Compounds of formula (I-B-6) can further be converted into a compound of formula (I-A), wherein R² is C₁₋₆alkylthio, said compound being represented by formula (I-A-16), by reaction with a suitable C₁₋₆alkylhalide, such as for example C₁₋₆alkyliodide, in the presence of a suitable base, such as dipotassium carbonate, and a suitable solvent, such as for example acetone.



- 10 Compounds of formula (I_{A/B}-a) can be converted into a compounds of formula (I-A) or (I-B), wherein X is N-R⁷, said compound being represented by formula (I_{A/B}-b), by reaction with an intermediate of formula (XVI), optionally in the presence of a suitable base, such as for example *N,N*-diethylethanamine, and in the presence of a suitable reaction-inert solvent, such as an alcohol, e.g. ethanol.

15



As already indicated in the preparation procedure of compounds of formula (I-A-13) described above, the compounds of formula (I) may also be converted to the


- 20 corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium
- 25 peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *tert*-butyl hydroperoxide. Suitable solvents are, for example, water, lower alkanols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones,
- 30 e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

Some of the intermediates and starting materials used in the above reaction procedures are commercially available, or may be synthesized according to procedures already described in the literature.

5


Intermediates of formula (II) may be prepared by reacting an intermediate of formula (XVII) with an intermediate of formula (XVIII), wherein W_5 represents a suitable leaving group such as a halo atom, e.g. chloro, bromo and the like, in the presence of magnesium, diethylether and a suitable reaction-inert solvent, such as diethylether.

10




Intermediates of formula (XVII) may be prepared by oxidizing an intermediate of formula (XIX) in the presence of a suitable oxidizing agent, such as MnO_2 , and a suitable reaction-inert solvent, such as methylene chloride.

15



Intermediates of formula (XIX) can be prepared by reducing an intermediate of formula (XX) in the presence of a suitable reducing agent such as lithium aluminium hydride, and a suitable reaction-inert solvent, such as tetrahydrofuran.


20



Intermediates of formula (XX), wherein Q represents a quinoline moiety optionally substituted in position 3 with C_{1-6} alkyl and wherein the carbonyl moiety is placed in position 6, said intermediates being represented by formula (XX-a), can be prepared by reacting an intermediate of formula (XXI) with an intermediate of formula (XXII) in

25


the presence of sodium 3-nitro-benzene sulfonate, a suitable acid, such as sulfuric acid, and a suitable alcohol, e.g. methanol, ethanol, propanol, butanol and the like.



5


Alternatively, intermediates of formula (II) can also be prepared by reacting an intermediate of formula (XXIII) with an intermediate of formula (XXIV), wherein W_6 is a suitable leaving group, such as a halo atom, e.g. bromo, chloro and the like, in the presence of a suitable agent, such as butyl lithium and a suitable reaction-inert solvent, such as tetrahydrofuran.

10




Intermediates of formula (XXIII) can be prepared by oxidizing an intermediate of formula (XXV) using the Moffatt Pfitzner or Swern oxidation (dimethylsulfoxide adducts with dehydrating agents e.g. DCC, Ac_2O , SO_3 , P_4O_{10} , COCl_2 or Cl-CO-COCl) in an inert solvent such as methylene chloride.

15




20

Intermediates of formula (XXV) can be prepared by reducing an intermediate of formula (XXVI) in the presence of a suitable reducing agent, such as for example lithium aluminium hydride and a suitable reaction-inert solvent, such as benzene.



25


Intermediates of formula (XXVI) can be prepared from an intermediate of formula (XXVII) by esterification in the presence of a suitable alcohol, such as methanol, ethanol, propanol, butanol and he like, and a suitable acid, such as sulfuric acid.



5


Intermediates of formula (XXVII), wherein R^1 represents a radical of formula (a-1) with Z_1 being O, Z_2 being CH_2 and n being 1, said intermediates being represented by formula (XXVII-a), can be prepared by reducing an intermediate of formula (XXVIII) in the presence of a suitable reducing agent such as hydrogen, and a suitable catalyst, such as palladium on charcoal, and a suitable acid such as acetic acid. When R^1 of intermediate (XXVII) represents an optionally substituted phenyl moiety, it can also be converted into an optionally substituted cyclohexyl moiety by reduction in the presence of a suitable reducing agent such as rhodium on Al_2O_3 , and a suitable reaction-inert solvent, such as tetrahydrofuran.

15




Intermediates of formula (IV), wherein Q represents a quinoline moiety substituted in position 2 with halo, e.g. chloro, said intermediates being represented by formula (IV-a), can be prepared by reacting an intermediate of formula (IV), wherein Q represents a quinolinone moiety with R^5 being hydrogen, said intermediate being represented by formula (IV-b), in the presence of POCl_3 .

20




Intermediates of formula (IV-a), wherein R^4 is hydrogen, said intermediates being represented by formula (IV-a-1), can also be prepared by reacting an intermediate of formula (XXIX) with POCl_3 in the presence of *N,N*-dimethylformamide (Vilsmeier-Haack formylation followed by cyclization).

5




Intermediates of formula (XXIX) may be prepared by reacting an intermediate of formula (XXX) with an intermediate of formula (XXXI), wherein W_7 represents a suitable leaving group, such as a halo atom, e.g. chloro, in the presence of a suitable base, such as for example *N,N*-diethylethanamine, and a suitable reaction-inert solvent, such as methylene chloride.

10



15 Intermediates of formula (IV-a) can be converted into an intermediate of formula (IV-c) by reaction with an intermediate of formula (XXXII) in the presence of a suitable reaction-inert solvent, such as an alcohol, e.g. methanol and the like.




20

Intermediates of formula (IV-a) can also be converted into an intermediate of formula (IV-d-1) by reaction with a suitable amine of formula (XXXIII-a), wherein Z_3 and Z_4 each independently represent hydrogen, C_{1-6} alkyl, C_{1-6} alkyloxy C_{1-6} alkyl, C_{1-6} alkylthio C_{1-6} alkyl or into an intermediate of formula (IV-d-2) by reaction with a suitable amine of formula (XXXIII-b), wherein Z_3 and Z_4 are taken together to form a


25

heterocycle as defined hereinabove in the definition of R^2 provided that the heterocycle comprises at least one nitrogen atom, in the presence of a suitable base, such as for example dipotassium carbonate, and a reaction-inert solvent, such as *N,N*-dimethylformamide.




5

Intermediates of formula (IV-a), wherein R^3 represents $\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-Cl}$, said intermediates being represented by formula (IV-a-2), can also be converted into an intermediate of formula (IV), wherein R^2 and R^3 are taken together to form a bivalent radical of formula $-\text{O-CH}_2\text{-CH}_2\text{-CH}_2-$, said intermediate being represented by formula (IV-e-1), by reaction with a suitable acid, such as hydrochloric acid and the like.
 Intermediates of formula (IV-a-2) can also be converted into an intermediate of formula (IV), wherein R^2 and R^3 are taken together to form a bivalent radical of formula $-\text{S-CH}_2\text{-CH}_2\text{-CH}_2-$, said intermediate being represented by formula (IV-e-2), by
 15 reaction with $\text{H}_2\text{N-C(=S)-NH}_2$ in the presence of a suitable reaction-inert solvent, such as an alcohol, e.g. ethanol.




Intermediates of formula (V) may be prepared by reacting an intermediate of formula (XXVII) with an intermediate of formula $\text{CH}_3\text{-NH-O-CH}_3$ in the presence of 1,1'-carbonyldiimidazole and a suitable reaction-inert solvent, such as methylene chloride.

5




Intermediates of formula (VII), wherein Q represents a quinoline moiety, in particular a quinoline moiety wherein R^2 is ethyl, R^3 is methyl and R^4 is hydrogen, and the carboxyl moiety is placed in position 6, said intermediates being represented by formula (VII-a),
10 can be prepared by reaction an intermediate of formula (XXXIV) in the presence of a suitable aldehyde, such as $\text{CH}_3\text{-CH}_2\text{-CH(=O)}$, $(\text{CH}_2\text{O})_n$, ZnCl_2 , FeCl_3 and a suitable reaction-inert solvent, such as an alcohol, for example ethanol.

15



Intermediates of formula (VIII) can be prepared by reacting an intermediate of formula (XXXV) with an intermediate of formula (XXXVI) in the presence of a suitable catalyst, such as for example tetrakis(triphenylphosphine)palladium and a suitable
20 reaction-inert solvent, such as for example dioxane.



25 Still some other preparations can be devised, some of them are disclosed further in this application with the Examples.

Pure stereoisomeric forms of the compounds and the intermediates of this invention may be obtained by the application of art-known procedures. Diastereomers may be separated by physical separation methods such as selective crystallization and chromatographic techniques, e.g. liquid chromatography using chiral stationary phases. Enantiomers may be separated from each other by the selective crystallization of their diastereomeric salts with optically active acids. Alternatively, enantiomers may be separated by chromatographic techniques using chiral stationary phases. Said pure stereoisomeric forms may also be derived from the corresponding pure stereoisomeric forms of the appropriate starting materials, provided that the reaction occurs stereoselectively or stereospecifically. Preferably, if a specific stereoisomer is desired, said compound will be synthesized by stereoselective or stereospecific methods of preparation. These methods will advantageously employ chirally pure starting materials. Stereoisomeric forms of the compounds of formula (I) are obviously intended to be included within the scope of the invention.

A stereoisomer of a compound of formula (I-A) or (I-B) such as a *cis* form, may be converted into another stereoisomer such as the corresponding *trans* form by reacting the compound with a suitable acid, such as hydrochloric acid, in the presence of a suitable reaction-inert solvent, such as for example tetrahydrofuran.

The mGluR1 antagonistic activity of the present compounds can be demonstrated in the Signal transduction on cloned rat mGluR1 in CHO cells test and the Cold allodynia test in rats with a Bennett ligation, as described hereinafter.

Due to their mGluR antagonistic activity, more in particular their Group I mGluR antagonistic activity and even more in particular, their mGluR1 antagonistic activity, the compounds of formula (I-A) or (I-B), their *N*-oxides, pharmaceutically acceptable addition salts, quaternary amines and stereochemically isomeric forms are useful in the treatment or prevention of glutamate-induced diseases of the central nervous system. Diseases in which a role for glutamate has been demonstrated include drug addiction or abstinence (dependence, opioid tolerance, opioid withdrawal), hypoxic, anoxic and ischemic injuries (ischemic stroke, cardiac arrest), pain (neuropathic pain, inflammatory pain, hyperalgesia), hypoglycemia, diseases related to neuronal damage, brain trauma, head trauma, spinal cord injury, myelopathy, dementia, anxiety, schizophrenia, depression, impaired cognition, amnesia, bipolar disorders, conduct disorders,

Alzheimer's disease, vascular dementia, mixed (Alzheimer's and vascular) dementia, Lewy Body disease, delirium or confusion, Parkinson's disease, Huntington's disease, Down syndrome, epilepsy, aging, Amyotrophic Lateral Sclerosis, multiple sclerosis, AIDS (Acquired Immune Deficiency Syndrome) and AIDS related complex (ARC).

5

The present invention also provides compositions for the administration to mammals, in particular humans, in particular for diagnostic reasons, more in particular for imaging an organ comprising a therapeutically effective amount of a radiolabelled compound of formula (I-A)* or (I-B)* and a pharmaceutically acceptable carrier or diluent.

10

Therefore, the compounds of the present invention or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, a therapeutically effective amount of a particular compound, in base or addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, rectally, topically, percutaneously or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, emulsions, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. As appropriate compositions for topical application there may be cited all compositions usually employed for topically administering drugs e.g. creams, gel, dressings, shampoos, tinctures, pastes,

35

ointments, salves, powders and the like. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause a significant deleterious effect to the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, suppositories, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

The diagnostically effective dose or frequency of administration depends on the particular compound of formula (I-A)* or (I-B)* used and the particular condition of the mammal being treated, as is well known to those skilled in the art.

The following examples are intended to illustrate and not to limit the scope of the present invention.


Experimental part

Hereinafter, "DMF" is defined as *N,N*-dimethylformamide, "DIPE" is defined as diisopropylether, "DMSO" is defined as dimethylsulfoxide, "BHT" is defined as 2,6-bis(1,1-dimethylethyl)-4-methylphenol, and "THF" is defined as tetrahydrofuran.

A. Preparation of the intermediates

Example A1

Preparation of



(interm. 1)

A mixture of 4-(1-methylethoxy)benzoic acid (0.083 mol) and Rh/Al₂O₃ 5% (10g) in THF (220ml) was hydrogenated at 50°C (under 3 bar pressure of H₂) for 1 night. The

mixture was filtered over celite, washed with THF and evaporated. Yield: 16g of intermediate 1 (100%).


Example A2

Preparation of 2-ethyl-3-methyl-6-quinolinecarboxylic acid (interm. 2)

- 5 A mixture of 4-aminobenzoic acid (0.299 mol) in ethanol (250ml) was stirred at room temperature. ZnCl_2 (0.0367 mol) and $(\text{CH}_2\text{O})_n$ (10g) were added. $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (0.5 mol) was added portionwise and the temperature rised till 60-65°C. Propanal (30ml) was added dropwise over a 2 hours period. The mixture was refluxed for 2 hours and kept at room temperature for 12 hours. The mixture was poured into water and filtered
10 through celite. The filtrate was acidified till pH=7 with HCl 6N and the mixture was evaporated till dryness. The residue was used without further purification. Yield : 56.1g of 2-ethyl-3-methyl-6-quinolinecarboxylic acid (interm. 2).

Example A3

Preparation of




(interm. 3)

- Pentanoyl chloride (0.2784 mol) was added at 5°C to a mixture of
15 4-bromobenzenamine (0.232 mol) and *N,N*-diethylethanamine (0.2784 mol) in CH_2Cl_2 (400ml). The mixture was stirred at room temperature overnight, poured out into water and extracted with CH_2Cl_2 . The organic layer was separated, washed with a concentrated NH_4OH solution and water, dried (MgSO_4), filtered and the solvent was evaporated. The residue (60g) was crystallized from diethylether. The precipitate was
20 filtered off and dried. The residue (35g, 63%) was taken up in CH_2Cl_2 . The organic layer was separated, washed with a 10% K_2CO_3 solution, washed with water, dried (MgSO_4), filtered and the solvent was evaporated. Yield: 30g of intermediate (3) (54%).

Example A4

Preparation of




(interm. 4)

A mixture of 6-bromo-2(1*H*)-quinolinone (0.089 mol) in POCl₃ (55ml) was stirred at 60°C overnight, then at 100°C for 3 hours and the solvent was evaporated. The residue was taken up in CH₂Cl₂, poured out into ice water, basified with NH₄OH conc., filtered over celite and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. Yield: 14.5g of intermediate (4) (67%).

Example A5

a) Preparation of




(interm. 5)

DMF (37ml) was added dropwise at 10°C under N₂ flow to POCl₃ (108ml). After complete addition, the mixture was allowed to warm to room temperature.

N-(4-bromophenyl)butanamide (0.33 mol) was added portionwise. The mixture was stirred at 85°C overnight, then allowed to cool to room temperature and poured out on ice (exothermic reaction). The precipitate was filtered off, washed with a small amount of water and dried (vacuum). The residue was washed with EtOAc/diethyl ether and dried. Yield: 44.2g of intermediate (5) (50%).

b) Preparation of




(interm. 6)

A mixture of intermediate (5) (0.162 mol) in methanol (600ml), and a solution of methanol sodium salt in methanol at 35% (154ml) was stirred and refluxed overnight. The mixture was poured out on ice. The precipitate was filtered off, washed with a small amount of water and taken up in CH₂Cl₂. K₂CO₃ 10% was added and the mixture was extracted with CH₂Cl₂. The organic layer was separated, washed with water, dried (MgSO₄), filtered and the solvent was evaporated. Yield: 31.9g of intermediate (6) (74%).

Example A6

Preparation of



(interm. 7)

1,1'-Carbonylbis-1*H*-imidazole (0.074 mol) was added portionwise to a mixture of 4-methoxycyclohexanecarboxylic acid (0.063 mol) in CH₂Cl₂ (200ml). The mixture


was stirred at room temperature for 1 hour. Then *N*-methoxymethanamine (0.074 mol) was added. The mixture was stirred at room temperature overnight, poured out into H₂O and extracted with CH₂Cl₂. The organic layer was separated, washed several times with H₂O, dried (MgSO₄), filtered and the solvent was evaporated. Yield: 12.6g of interm. 7.

Example A7

- a) A mixture of 6-fluoro-4-oxo-4*H*-1-benzopyran-2-carboxylic acid (0.30mol) in acetic acid (400ml) was hydrogenated with Pd/C (3g) as a catalyst. After uptake of H₂ (3 equiv), the catalyst was filtered off. The filtrate was evaporated. The residue was stirred in petroleum ether. The precipitate was filtered off and dried (vacuum; 70°C). After recrystallization from CHCl₃/CH₃OH, the precipitate was filtered off and dried (vacuum; 80°C and high vacuum; 85°C). Yield: 8.8 g of 6-fluoro-3,4-dihydro-2*H*-1-benzopyran-2-carboxylic acid (interm. 8) (15.0%).
- b) A mixture of intermediate (8) (0.255 mol) in ethanol (400ml) and H₂SO₄ (5ml) was stirred and refluxed for 8 hours. The solvent was evaporated till dryness. The residue was dissolved in CH₂Cl₂. The organic layer was separated, washed with K₂CO₃ 10%, dried (MgSO₄), filtered and the solvent was evaporated. Yield: 45g of ethyl 6-fluoro-3,4-dihydro-2*H*-1-benzopyran-2-carboxylate (interm. 9) (79%).
- c) Reaction under N₂. A mixture of sodium bis(2-methoxyethoxy)aluminumhydride, 70 wt % solution in methylbenzene 3.4M (0.44 mol) in benzene (150 ml) (reflux) was added dropwise during 1 hour to a refluxed mixture of interm. 9 (0.22 mol) and benzene (600 ml). After stirring for 2.5 hours at reflux temperature, the mixture was cooled to ±15°C. The mixture was decomposed by adding dropwise ethanol (30 ml) and water (10 ml). This mixture was poured out onto ice/water and this mixture was acidified with concentrated hydrochloric acid. This mixture was extracted with diethyl ether (500 ml). The separated organic layer was washed with water, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent : CHCl₃). The desired fraction was collected and the eluent was evaporated. Yield: 34 g of 6-fluoro-3,4-dihydro-2*H*-1-benzopyran-2-methanol (interm. 10) (85%).
- d) Reaction under N₂. To a stirred and cooled (-60°C; 2-propanone/CO₂ bath) mixture of ethanedioyl dichloride (0.1 mol) in CH₂Cl₂ (350 ml) was added sulfinylbis[methane] (30 ml) during 10 minutes. After stirring 10 minutes, a mixture of interm. 10 in CH₂Cl₂ (90 ml) was added during 5 minutes. After stirring for 15 minutes, *N,N*-diethylethanamine (125 ml) was added. When the mixture was warmed up to room temperature, it was poured out in water. The product was extracted with CH₂Cl₂.

The organic layer was washed with water, HCl (1M), water, NaHCO₃ (10%) and water, dried and evaporated. The residue was dissolved in diethyl ether, washed with water, dried, filtered and evaporated. The residue was purified by column chromatography over silica gel (eluent : CHCl₃). The desired fraction was collected and the eluent was evaporated. Yield: 21.6 g of 6-fluoro-3,4-dihydro-2*H*-1-benzopyran-2-carboxaldehyde (interm. 11) (67%).

e) Preparation of




(interm. 12)

nButyllithium 1.6M (0.056 mol) was added slowly at -70°C to a solution of intermediate (5) (0.046 mol) in THF (100ml). The mixture was stirred at -70°C for 30 minutes. A suspension of interm. 11 (0.056 mol) in THF (100ml) was added slowly.

The mixture was stirred at -70°C for 1 hour, then brought to room temperature, poured out into H₂O and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (21g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 80/10; 15-35μm). The pure fractions were collected and the solvent was evaporated. Yield: 9.5g of interm. 12 (55%).


Example A8

a) Preparation of



(interm. 13)

and




(interm. 14)

A mixture of intermediate (5) (0.1127 mol), 2-methoxyethanamine (0.2254 mol) and K₂CO₃ (0.2254 mol) in DMF (500ml) was stirred at 120°C for 15 hours and then cooled. The solvent was evaporated. The residue was taken up in CH₂Cl₂ and H₂O.

The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated till dryness. The residue (33.53g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 99.5/0.5; 15-40 μm). Two fractions were collected and their solvents were evaporated. Yield: 5.7g of interm. 14 (38%) and interm. 13 (34%).

b) Preparation of



(interm. 15)

A mixture of intermediate (5) (0.0751 mol), thiomorpholine (0.0891 mol) and K₂CO₃ (0.15 mol) in DMF (200ml) was stirred at 120°C for 12 hours. The solvent was


evaporated till dryness. The residue was taken up in CH_2Cl_2 and H_2O . The organic layer was separated, dried (MgSO_4), filtered and the solvent was evaporated. The residue (26g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 80/20; 20-45 μm). Two fractions were collected and their solvents were evaporated. The two fractions were combined. Yield: 9.4g of interm. 15 (37%); mp. 82°C.

Example A9

- a) 4-Aminobenzoic acid (0.219 mol) was added to a solution of sodium 3-nitrobenzenesulfonate (0.118 mol) in H_2SO_4 70% (230ml) and the mixture was stirred and refluxed. 2-propene-1,1-diol, 2-methyl-, diacetate (0.216 mol) was added dropwise and the mixture was refluxed for 4 hours. Ethanol (200ml) was added and the mixture was stirred at 80°C for 48 hours. The mixture was evaporated, the residue was poured into ice water/ NH_4OH and extracted with CH_2Cl_2 . The organic layer was dried (MgSO_4) and evaporated. The residue was purified by column chromatography over silica gel (eluent : CH_2Cl_2 /2-propanol 99/1). The pure fractions were collected and evaporated. Yield : 21g of ethyl 3-methyl-6-quinolinecarboxylate (interm. 16) (45%).
- b) Interm. 16 (0.098 mol) in THF (270ml) was added at 0°C to a solution of LiAlH_4 (0.098 mol) in THF under N_2 . When the addition was complete, water (10ml) was added. The precipitate was filtered off and washed with CH_2Cl_2 . The organic layer was dried (MgSO_4), filtered off and evaporated. The product was used without further purification. Yield : 16.71g of 3-methyl-6-quinolinemethanol (interm. 17).
- c) MnO_2 (0.237 mol) was added to a solution of interm. 17 (0.096 mol) in CH_2Cl_2 (200ml) and the mixture was stirred at room temperature for 12 hours. The mixture was filtered through celite and the filtrate was stirred again with MnO_2 (20g) for 12 hours. MnO_2 (10g) was added again. The mixture was stirred for 12 hours. The mixture was filtered through celite and evaporated. The product was used without further purification. Yield : 11.71g of 3-methyl-6-quinolinecarboxaldehyde (71%) (interm. 18).
- d) A solution of bromocyclohexyl (0.14 mol) in 1,1'-oxybisethane (50ml) and Mg turnings (50ml) was added at 10°C to a mixture of THF (0.14 mol) in 1,1'-oxybisethane (10ml). A solution of interm. 18 (0.07 mol) in Mg turnings (100ml) was added carefully at 5°C, the mixture was poured into ice water and extracted with EtOAc. Yield : 11.34g of (\pm)- α -cyclohexyl-3-methyl-6-quinolinemethanol (63%) (interm. 19).

Example A10


Preparation of



A mixture of compound (5) (0.001507 mol), tributyl(1-ethoxyethenyl)stannane (0.00226 mol) and Pd(PPh₃)₄ (0.000151 mol) in 1,4-dioxane (5ml) was stirred at 80°C for 3 hours. Water was added. The mixture was extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. This product was used without further purification. Yield: 1.4g of interm. 20.

Example A11



Preparation of



A mixture of compound (45) (prepared according to B6) (0.00125 mol) in NaOH 3N (5 ml) and iPrOH (1.7 ml) was stirred at room temperature overnight, then poured out into H₂O, acidified with HCl 3N and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue was taken up in diethyl ether. The precipitate was filtered off and dried. Yielding: 0.26 g of intermediate 21 (56%). (mp.: 232°C)


Example A12

a. Preparation of




A mixture of 5-bromo-1H-indole-2,3-dione (0.221 mol) in NaOH 3N (500 ml) was stirred at 80°C for 30 minutes, brought to room temperature and 2-pentanone (0.221 mol) was added. The mixture was stirred and refluxed for 1 hour and 30 minutes and acidified with AcOH until pH=5. The precipitate was filtered, washed with water and dried. Yielding 52.3 g of intermediate 22 and intermediate 23. (Total yielding: 80%).

b. Preparation of



(interm. 24)




(interm. 25)

nBuLi 1.6 M (0.0816 mol) was added dropwise at -78°C to a suspension of intermediate 22 (0.034 mol) and intermediate 23 (0.034 mol) in THF (300 ml) under N_2 flow. The mixture was stirred at -78°C for 30 minutes. nBuLi 1.6M (0.0816 mol) was added dropwise. The mixture was stirred for 1 hour. A mixture of intermediate 9 (0.102 mol) in THF (250 ml) was added slowly. The mixture was stirred for -78°C to -20°C , poured out into $\text{H}_2\text{O}/\text{HCl}$ 3N and extracted with EtOAc. The organic layer was separated, dried (MgSO_4), filtered, and the solvent was evaporated till dryness. Yielding: 20.89 g of compound intermediate 24 and intermediate 25 (86%).

Example A13


a. Preparation of



(interm. 26)

4-amino-3-methoxybenzoic acid (0.054 mol) was added portionwise at room temperature to a solution of 3-chloro-2-ethyl-2-butenal (0.065 mol) in AcOH (100ml). The mixture was stirred and refluxed for 8 hours and evaporated to dryness. The residue was taken up in CH_2Cl_2 , water was added and the solution was basified by Et_3N . The organic layer was separated, dried (MgSO_4), filtered, and the solvent was evaporated. The residue was crystallized from 2-propanone. The precipitate was filtered off and dried. Yielding: 2.5g of interm. 26 (18%).

b. Preparation of




(interm. 27)

CDI (0.012 mol) was added at room temperature to a solution of interm. 26 (0.011 mol) in CH_2Cl_2 (30ml). The mixture was stirred at room temperature for 1 hour. methoxyaminomethyl (0.012 mol) was added and the mixture was stirred at room temperature for 8 hours. H_2O was added. A precipitate was filtered off. The filtrate was extracted with CH_2Cl_2 . The organic layer was separated, dried (MgSO_4), filtered, and

the solvent was evaporated. The residue was crystallized from diethyl ether. The precipitate was filtered off and dried. Yielding: 0.95g of interm. 27 (31%) (mp.:148°C).

Example A14




Preparation of

(interm. 28)

- 5 4-Bromobenzenamine (0.034 mol) was added at room temperature to a solution of 3-chloride-2-ethyl-2-butanal (0.041 mol) in AcOH (60 ml). The mixture was stirred and refluxed for 8 hours, brought to room temperature and evaporated to dryness. The product was crystallized from EtOAc. The precipitate was filtered, washed with K_2CO_3 10% and taken up in CH_2Cl_2 . The organic layer was separated, dried ($MgSO_4$),
10 filtered, and the solvent was evaporated. Yielding: 4,6 g of interm. 28 (54%).

Example A15


a. Preparation of



(interm. 29)


- A solution of KOH (0.326 mol) in H_2O (150ml) was added slowly at 5°C to a solution of 1,3-cyclohexanedione (0.268 mol) in H_2O (150ml). The temperature must not reach 12 °C. KI (2g) then 2-bromo-1-(4-nitrophenyl)ethanone (0.294 mol) were added
15 portionwise. The mixture was stirred at room temperature for 48 hours. The precipitate was filtered, washed with H_2O then with diethyl ether and dried. Yielding: 63g (85%). A part of this fraction (1g) was crystallized from EtOH. The precipitate was filtered off and dried. Yielding: 0.5g of interm. 29 (42%) (mp.: 100°C).

b. Preparation of




(interm. 30)

- A mixture of interm. 29 (0.145 mol) in H_2SO_4 (40ml) was stirred at room temperature
20 for 1 hour, poured out into ice, basified with NH_4OH and extracted with CH_2Cl_2 . The organic layer was separated, dried ($MgSO_4$), filtered, and the solvent was evaporated. The residue was crystallized from EtOH. The precipitate was filtered off and dried. Yielding: 31g (58%). A part of this fraction (1g) was crystallized from EtOH. The precipitate was filtered off and dried. Yielding: 0.7g of interm. 30 (58%) (mp.:200°C).




A mixture of interm. 30 (0.039 mol), Raney Ni (10g) in EtOH (100ml) was hydrogenated at room temperature under a 3 bar pressure for 1 hour. The mixture was filtered over celite and washed with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue (9.5g) was crystallized from diethyl ether. The precipitate was filtered off and dried. Yielding: 4.6g (52%). The filtrate was evaporated. The residue (2.7g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH; 99/1; 15-40μm). Two fractions were collected and the solvent was evaporated. Yielding: 1.6g F1 and 1.2g F2. F2 was crystallized from EtOH. The precipitate was filtered off and dried. Yielding: 0.24g of interm. 31 (2%) (mp.: 202°C).



Interm. 30 (0.02 mol) was added at room temperature to a solution of 3-chloro-2-ethyl-2-butenal (0.04 mol) in AcOH (50ml). The mixture was stirred and refluxed for 4 hours. The solvent was evaporated till dryness. The residue was crystallized from EtOAc. The precipitate was filtered off and dried. The residue was taken up in CH₂Cl₂. The mixture was basified with K₂CO₃ 10% and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue was crystallized from EtOH. The precipitate was filtered off and dried. Yielding: 2.5g of interm. 32 (40%).


Example A16



A mixture of 2-(4-nitrophenyl)-1-phenylethanone (0.083 mol) and Raney Ni (20g) in EtOH (200ml) was hydrogenated at room temperature for 1 hour under a 3 bar pressure, then filtered over celite, washed with CH₂Cl₂/CH₃OH and dried. Yielding: 17.5g of interm. 33 (97%).

Example A17


a. Preparation of



(interm. 34)

DMF (12.4ml) was added dropwise at 5°C to POCl₃ (0.7536 mol). 4'-bromo-5-chlorovaleranilide (0.1032 mol) was added and the mixture was stirred at 75°C for 6 hours, cooled at room temperature and poured out into ice water. The insoluble was
5 filtered, washed with water and dried. Yielding: 25.7g of intermediate 34 (78%).

b. Preparation of




(interm. 35)

A mixture of intermediate 34 (0.094 mol) in HCl 6N (250ml) was stirred and refluxed for 2 days, cooled, poured out on water (100ml) and neutralized with NH₄OH (concentrated). The insoluble was filtered and washed with water then with EtOH. Yielding: 19g. The filtrate was evaporated. The residue (9.4g) was purified by column
10 chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 99.25/0.75; 15-35μm). One fraction was collected and the solvent was evaporated. Yielding: 8g of intermediate 35 (32%).

15 B. Preparation of the non-radioactive compounds

Example B1

Preparation of




(compound 306)

POCl₃ (0.024 mol) was added slowly at 5°C to DMF (0.024 mol). The mixture was stirred at room temperature for 30 minutes, then cooled to 5°C. 3-Oxo-butanoic acid ethyl ester (0.024 mol) was added slowly. The mixture was stirred at 5°C for 30
20 minutes. 1-(4-aminophenyl)-2-phenylethanone (0.024 mol) was added portionwise. The mixture was stirred at 90°C for 3 hours and dissolved in CH₂Cl₂. Ice water was added. The mixture was basified with NH₄OH and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue was crystallized from 2-propanone/diethyl ether. The precipitate was
25 filtered off and dried. Yielding: 0.9 g of compound 306 (11%) (mp.:136°C).


Example B2

Preparation of



(compound 2)

KMnO₄ (10g) was added portionwise at room temperature to a solution of




(prepared according to example A7.e) (0.022 mol)

in tris(dioxa-3,6-heptyl)amine (1ml) and CH₂Cl₂ (100ml). The mixture was stirred at room temperature for 8 hours, filtered over celite, washed with CH₂Cl₂ and dried. The residue (6g, 100%) was crystallized from diethyl ether/petroleum ether. The precipitate was filtered off and dried. Yield: 2g of compound (2) (33%); mp. 82°C.

Example B3


a) Preparation of



(compound 3)

nBuLi 1.6M (0.07 mol) was added slowly at -70°C to a solution of intermediate (5) (0.058 mol) in THF (150ml). The mixture was stirred at -70°C for 30 minutes. A solution of 2,3-dihydro-1*H*-Indene-2-carbonitrile (0.07 mol) in THF (100ml) was added slowly. The mixture was stirred at -70°C for 1 hour, brought slowly to room temperature, hydrolyzed with H₂O and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (22g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/ cyclohexane 80/20 to 100; 15-35μm). The pure fractions were collected and the solvent was evaporated. The second fraction was crystallized from 2-propanone/diethyl ether. The precipitate was filtered off and dried. Yield: 0.11g of compound (3). The filtrate was concentrated. Yield: 0.55g of compound (3); mp. 145°C.

b) Preparation of




cis (compound 4)


trans (compound 5)

nBuLi 1.6M (0.022 mol) was added slowly at -70°C to a solution of intermediate (5) (0.018 mol) in THF (50ml). The mixture was stirred at -70°C for 1 hour, brought to -40°C, then cooled to -70°C. A solution of interm. 7 (0.018 mol) in THF (40ml) was added slowly. The mixture was stirred at -70°C for 1 hour, then brought to -20°C, hydrolyzed with H₂O and extracted with EtOAc. The organic layer was separated,

dried (MgSO_4), filtered and the solvent was evaporated. The residue (6.5g) was purified by column chromatography over silica gel (eluent: toluene/EtOAc 90/10; 15-40 μM). Two fractions (F1 and F2) were collected and the solvent was evaporated. F1 (2.4g) was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 1.8g of compound (4) (29%); mp. 123°C. F2 (0.9g) was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.2g of compound (5) (3%); mp. 120°C.



nBuLi 1.6M in exane (0.107 mol) was added dropwise at -78°C under N_2 flow to a mixture of intermediate (6) (0.089 mol) in THF. The mixture was stirred at -78°C for 1 hour. A mixture of interm. 7 (150 ml) was added at -78°C under N_2 flow. The mixture was stirred at -78°C for 2 hours, brought to 0°C, poured out into H_2O and extracted with EtOAc. The organic layer was separated, dried (MgSO_4), filtered and the solvent was evaporated. The residue (31g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 85/15; 20-45 μm). Two pure fractions were collected and their solvents were evaporated. Yielding: 11 g of compound (7) (38%) and 8.2 g of compound (8) (28%).




A solution of chloromethylbenzene (0.0069 mol) in diethyl ether (8ml) was added slowly to a suspension of Mg (0.0069 mol) in a small amount of diethyl ether. The mixture was stirred at room temperature for 30 minutes (disappearance of Mg), then cooled to 5°C. A solution of interm. 27 (0.0027 mol) in THF (8ml) was added slowly. The mixture was stirred at 5°C for 15 minutes, then at room temperature for 2 hours, poured out into H_2O and filtered over celite. The precipitate was washed with EtOAc. The filtrate was extracted with EtOAc. The organic layer was separated, dried (MgSO_4), filtered, and the solvent was evaporated. The residue (1g) was purified by column chromatography over kromasil (eluent: CH_2Cl_2 100 to $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 99/1; 15-40 μm). The pure fractions were collected and the solvent was evaporated. The

residue (0.5g, 56%) was crystallized from diethyl ether. The precipitate was filtered off and dried. Yielding: 0.14g of compound 503 (15%).


Example B4: examples of endgroup modifications

a) Preparation of



(compound 156)


A mixture of



(compound 8)


- 5 (prepared according to example B3.c) (0.018 mol) in HCl 3N (60ml) and THF (60ml) was stirred at 60°C overnight. The mixture was basified with a K₂CO₃ 10% solution and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. Yield: 4.6g of compound (156) (82%).

b) Preparation of



(compound 9)



A mixture of




(compound 7)

- 10 (prepared according to example B3.c) (0.0122 mol) in HCl 3N (40ml) and THF (40ml) was stirred and refluxed overnight, poured out into water, basified with K₂CO₃ 10% and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 40/60; 15-40μm). The pure fractions were
- 15 collected and the solvent was evaporated. Yield: 2g of compound (9) (52%); mp. 226°C.


c) Preparation of



A mixture of compound (4) (0.0015 mol), 2-methoxyethanamine (0.003 mol) and K_2CO_3 (0.003 mol) in DMF (5ml) was stirred at 140°C for 48 hours. H_2O was added. The mixture was extracted with EtOAc. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (1g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 60/40; 15-40 μ m). Two fractions were collected and the solvent was evaporated. Both fractions were crystallized separately from pentane. The precipitate was filtered off and dried. Yield: 0.05g of compound (10) (9%; mp. 115°C) and 0.057g of compound (11) (10%; mp. 107°C).




A mixture of compound (4) (0.0015 mol) in 2-(methylthio)ethanamine (2ml) was stirred at 120°C for 8 hours. K_2CO_3 10% was added. The mixture was extracted with CH_2Cl_2 . The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (2.2g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 70/30; 15-40 μ m). Two fractions were collected and the solvent was evaporated. The first fraction was crystallized from diethyl ether/petroleum ether. The precipitate was filtered off and dried. Yield: 0.08g of compound (12) (14%); mp. 120°C. The second fraction was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.18g of compound (13) (31%); mp. 125°C.



A mixture of compound (4) (0.001507 mol), ethynyltrimethylsilane (0.003013 mol), CuI (0.000151 mol) and $Pd(PPh_3)_4$ (0.000151 mol) in *N,N*-diethylethanamine (5ml) was stirred at 100°C for 24 hours. Water was added. The mixture was filtered over celite, washed with EtOAc and the filtrate was extracted with EtOAc. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (1.3g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 85/15; 15-40 μ m). The pure fractions were collected and the solvent was evaporated. The residue (0.3g) was crystallized from pentane. The precipitate was filtered off and dried. Yield: 0.11g of compound (14) (18%); mp. 114°C.

f) Preparation of




(compound 15)


cis

A mixture of compound (14) (0.013 mol) and KF (0.038 mol) in acetic acid (50ml) was stirred at room temperature for 2 hours. H₂O was added and the mixture was extracted with diethyl ether. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (4.4g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 70/30; 15-40μm). One fraction was collected and the solvent was evaporated. This fraction (3g, 73%) was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 2.45g of compound (15) (60%); mp. 132°C.

g) Preparation of




cis (compound 15) and




trans (compound 17)

A mixture of



cis (compound 14) and




trans (compound 16)

prepared according to example B.7.a) (0.0056 mol) in KOH [1M, H₂O] (10ml) and methanol (30ml) was stirred at room temperature for 1 hour, poured out into water and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (2.2g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 85/15 to 70/30; 15-40μm). Two fractions were collected and the solvent was evaporated.

The first fraction was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.2g of compound (15) (11%); mp. 133°C.

The second fraction was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.3g of compound (17) (16%); mp. 128°C.

h) Preparation of




(compound 18)

cis

A mixture of compound (4) (0.001205 mol), 2-propyn-1-ol (0.002411 mol), Pd(PPh₃)₄ (0.000121 mol) and CuI (0.000121 mol) in *N,N*-diethylethanamine (5ml) was stirred at 100°C for 2 hours. Water was added. The mixture was filtered over celite, washed with EtOAc and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (0.7g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 98/2; 15-40 μm). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from petroleum ether and diethyl ether. The precipitate was filtered off and dried. Yield: 0.1g of compound (18) (23%); mp. 113°C.

i) Preparation of




A mixture of compound (4) (0.006027 mol) and KF (0.024108 mol) in DMSO (20ml) was stirred at 140°C. The solvent was evaporated till dryness. The residue was solidified in water and diethyl ether. The mixture was extracted with diethyl ether. The organic layer was separated, washed with diethyl ether, washed with a saturated solution of NaCl, dried (MgSO₄), filtered and the solvent was evaporated. The residue (1.7g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 85/15; 15-40 μm). Three fractions were collected and their solvents were evaporated.

The first fraction was crystallized from petroleum ether. The precipitate was filtered off and dried. Yield: 0.21g of compound (19) (11%); mp. 92°C.


The second fraction was crystallized from petroleum ether. The precipitate was filtered off and dried. Yield: 0.33g of compound (20) (17%); mp. 114°C.

j) Preparation of




A mixture of compound (4) (0.003013 mol), acetyl chloride (0.003315 mol) and sodium iodide (0.006027 mol) in CH₃CN (10ml) was stirred and refluxed for 1 hour. K₂CO₃ 10% was added. The mixture was extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (1g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 80/20; 15-40 μm). Two fractions were collected and their solvents

were evaporated. The first fraction was crystallized from petroleum ether. The precipitate was filtered off and dried. Yield: 0.12g of compound (21); mp. 110°C.




A mixture of compound (21) (0.000898 mol), trimethylsilanecarbonitrile (0.001347 mol) and Pd(PPh₃)₄ (0.00009 mol) in *N,N*-diethylethanamine (5ml) was stirred at 100°C for 2 hours. Water was added. The mixture was extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (0.4g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 80/20; 15-40 μm). The pure fractions were collected and the solvent was evaporated. The residue (0.18g, 62%) was crystallized from petroleum ether. The precipitate was filtered off and dried. Yield : 0.13g of compound (22) (45%); mp. 138°C.




A mixture of compound (4) (0.00603 mol), Pd(OAc)₂ (0.000603 mol), PPh₃ (0.00904 mol) and K₂CO₃ (0.012054 mol) in CO (gas) and methanol (40ml) was stirred at 90°C for 8 hours under a 5 bar pressure of CO. H₂O was added. The mixture was extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (6g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 100/0 to 98/2; 15-35 μm). Four fractions (F1-F4) were collected and the solvent was evaporated. Yield: 0.13g (cis) F1; 0.02g F2 (cis, compound 25); 0.055g F3 (trans, 3%) and 0.11g F4 (trans; compound 26). F1 was crystallized from petroleum ether. The precipitate was filtered off and dried. Yield: 0.03g of compound (23) (1%); mp. 91°C.


F3 was crystallized from petroleum ether. The precipitate was filtered off and dried. Yield: 0.035g of compound (24) (1%); mp. 99°C.



A mixture of compound (4) (0.009 mol) and Zn (0.027 mol) in acetic acid (30ml) was stirred at 60°C for 4 hours, filtered over celite, washed with CH₂Cl₂, evaporated till dryness, solubilized in CH₂Cl₂ and washed with K₂CO₃ 10%. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (4g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 75/25; 15-40µm). One fraction was collected and the solvent was evaporated. This fraction (1g 37%) was crystallized from petroleum ether. The precipitate was filtered off and dried. Yield: compound (25); mp. 88°C.




A mixture of compound (4) (0.001502 mol), Sn(CH₃)₄ (0.003004 mol) and Pd(PPh₃)₄ (0.00015 mol) in methylbenzene (5ml) was stirred and refluxed for 3 hours. K₂CO₃ 10% was added. The mixture was extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (0.7g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 85/15; 15-40 µm). Two fractions (F1 and F2) were collected and their solvents were evaporated. Yield: 0.27g (F 1, starting material) and 0.14g (F2). F2 was crystallized from pentane and petroleum ether. The precipitate was filtered off and dried. Yield: 0.08g of compound (27) (17%); mp. 110°C.



A mixture of compound (4) (0.001507 mol), tributylethenylstannane (0.002260 mol) and Pd(PPh₃)₄ (0.000151 mol) in dioxane (5ml) was stirred at 80°C for 8 hours. Water was added. The mixture was filtered over celite, washed with EtOAc and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (0.65g) was purified by column chromatography over

silica gel (eluent: cyclohexane/EtOAc 90/10; 15-40 μ m). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from petroleum ether. The precipitate was filtered off and dried. Yield: 0.07g of compound (28) (14%); mp. 108°C.

p) Preparation of




(compound 29)

trans

- 5 A mixture of compound (5) (0.001507 mol), triphenyl(phenylmethyl)stannane (0.002260 mol) and Pd(PPh₃)₄ (0.000151 mol) in dioxane (5ml) was stirred at 80°C for 8 hours. Water was added. The mixture was extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (1.4g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/EtOAc 96/4; 15-40 μ m). The pure fractions were collected and the solvent was evaporated.
- 10 The residue (0.38g) was crystallized from petroleum ether. The precipitate was filtered off and dried. Yield: 0.16g of compound (29) (28%); mp. 112°C.

q) Preparation of




(compound 30)

cis

- A mixture of compound (4) (0.001507 mol), tributyl-2-thienylstannane (0.00226 mol) and Pd(PPh₃)₄ (0.0001507 mol) in dioxane (5ml) was stirred at 80°C for 8 hours.
- 15 K₂CO₃ 10% was added. The mixture was extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (1.7g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 85/15; 15-40 μ m). The pure fractions were collected and the solvent was evaporated. The residue (0.65g) was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.35g of compound (30) (61%); mp. 142°C.
- 20

r) Preparation of




(compound 31)

cis

A mixture of compound (4) (0.0015 mol), 3-thienyl boronic acid (0.00226 mol), Pd(PPh₃)₄ (0.00015 mol) and dioxane was stirred and refluxed for 24 hours. K₂CO₃

5


s) Preparation of



cis


10

t) Preparation of



15


u) Preparation of



cis


A mixture of compound 4 (0.015 mol) and NaN_3 (0.045 mol) in DMF (50ml) was stirred at 140°C for 2 hours. K_2CO_3 10% was added and the mixture was extracted with EtOAc. The organic layer was separated, dried (MgSO_4), filtered and the solvent was evaporated. The residue (6g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 60/40; 15-40 μm). The first fraction was collected and the solvent was evaporated. The residue was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 1.26g of compound (35) (24%); mp. 160°C .

v) Preparation of



CIS

(compound 36)




TRANS

(compound 37)


A mixture of compound (4) (0.009 mol) and thiourea (0.0099 mol) in ethyl alcohol (30ml) was stirred and refluxed for 12 hours and a solution of KOH (0.0149 mol) in H_2O (5ml) was added slowly. The mixture was stirred and refluxed for 1 hour, poured out into water and extracted with CH_2Cl_2 . The organic layer was separated, dried (MgSO_4), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (cyclohexane/EtOAc 70/30; 15-40 μm). The pure fractions were collected and the solvent was evaporated. Yielding: 1.1g of F1 (37%) and 0.4g of F2 (13%). F1 was crystallized from 2-propanone. The precipitate was filtered off and dried. Yielding: compound (36). F2 was crystallized from 2-propanone. The precipitate was filtered off and dried. Yielding: compound (37).

w) Preparation of



CIS

(compound 38)




TRANS

(compound 39)

CH_3I (0.0034 mol) was added slowly at room temperature to a solution of compound (36) (0.0015 mol), compound (37) (0.0015 mol) and K_2CO_3 (0.0034 mol) in acetone (15ml). The mixture was stirred at room temperature for 8 hours. Water was added and the mixture was extracted with CH_2Cl_2 . The organic layer was separated, dried (MgSO_4), filtered and the solvent was evaporated. The residue (1.2g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 85/15; 15-40 μm). The pure fractions were collected and the solvent was evaporated. Yielding: 0.6g F1 (57%), and 0.18g F2 (17%). F1 was crystallized from diethyl ether. The precipitate was filtered off and dried. Yielding: 0.28g compound (38) (27%). F2 was crystallized from

diethyl ether. The precipitate was filtered off and dried. Yielding: 0.065g of compound (39) (6%).


x) Preparation of



(compound 40)

cis


A mixture of



compound (41) prepared

according to example B3.b (0.0014 mol) in HCl 3N (5ml) and THF (5ml) was stirred and refluxed for a weekend, then poured out into H₂O, basified with K₂CO₃ and
5 extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. Yielding: 0.5g of F. This fraction F was crystallized from 2-propanone. The precipitate was filtered off and dried. Yielding: 0.35g of compound (40) (74%).


y) Preparation of



(compound 188)

A mixture of compound (5) (0.045 mol), acetamide (0.90013 mol) and K₂CO₃ (0.225
10 mol) was stirred and refluxed at 200°C for 2 hours, cooled at room temperature, poured out into H₂O/CH₂Cl₂; and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated till dryness. The residue (14.4 g) was crystallized from CH₃OH. The precipitate was filtered off and dried. The filtrate was evaporated. The residue (11.27g) was purified by column chromatography
15 over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 96/4/0.1; 15-35μm). The pure fractions were collected and the solvent was evaporated. Yielding: 4.2 g of compound (188) (65%).


z) Preparation of



(compound 248)

A mixture of compound (188) (0.00032 mol), benzoic acid (1.5 equiv., 0.00048 mol), 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide .HCl (1:1) (1.5 equiv., 0.00048 mol), *N*-hydroxybenzotriazole (1.5 equiv., 0.00048 mol) and Et₃N (1 equiv., 0.00032 mol) in CH₂Cl₂ (2ml) was stirred at room temperature for 15 hours. The solvent was
5 evaporated. The residue was purified by HPLC and the product fractions were collected and the solvent was evaporated. Yield: 0.066 g of compound (205) (49.50%).

aa) Preparation of




(compound 6)

trans

A mixture of interm. 20 (0.001507 mol) in HCl 3N (10ml) and THF (10ml) was stirred at room temperature for 8 hours, basified with K₂CO₃ 10% and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was
10 evaporated. The residue (1.2g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 85/15; 15-40 μm). The pure fractions were collected and the solvent was evaporated. The residue (0.4g) was crystallized from petroleum ether. The precipitate was filtered off and dried. Yield: 0.3g of compound (6) (58%); mp. 108°C.

ab) Preparation of



(compound 419)


cis

A mixture of compound 213 (prepared according to B4) (0.00305 mol) and CH₃ONa (30% in CH₃OH) (0.00916 mol) in CH₃OH (25ml) was stirred and refluxed for 15 hours then cooled to room temperature, poured out into H₂O and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was
20 evaporated till dryness. The residue (1.1g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc; 40/60; 15-40μm). Two fractions were collected and the solvent was evaporated. Yielding: 0.3g F1 and 0.5g F2 (50%) F2 was crystallized from diethyl ether/petroleum ether. The precipitate was filtered off and dried. Yielding: 0.26g F1 was crystallized from pentane. The precipitate was filtered off and dried. Yielding: 0.19g. This fraction was purified by column chromatography
25 over silica gel (eluent: CH₂Cl₂/CH₃OH; 98/2; 15-40μm). The pure fractions were collected and the solvent was evaporated. Yielding: 0.11g. This fraction was purified by column chromatography over kromasil (eluent: CH₃OH/H₂O; 70/30). The pure


fractions were collected and the solvent was evaporated. Yielding: 0.09g. (9%) This fraction was crystallized from diethyl ether. The precipitate was filtered off and dried. Yielding: 0.08g of compound 419 (8%).

Example B5

Preparation of



cis (compound 42)




(trans) (compound 43)


- 5 Iodomethane (0.00456 mol) was added at 5°C to a mixture of compound (9) (0.0019 mol), compound (8) (0.0019 mol) and tBuOK (0.00456 mol) in THF (30ml) under N₂ flow. The mixture was stirred at room temperature overnight, poured out into H₂O and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue was purified by column chromatography over
10 silica gel (eluent: cyclohexane/EtOAc 65/35; 15-40μm). Two fractions were collected and the solvent was evaporated. Yield: 0.35g of compound (42) (30%; mp. 125°C) and 0.35g of compound (43) (30%; mp. 116°C).

Example B6

a) Preparation of




cis (compound 44)




(trans) (compound 45)

- NaH 60% (0.01068 mol) was added at 0°C under N₂ flow to a mixture of compound (8) and compound (9) (0.0089 mol). The mixture was stirred for 30 minutes.
Ethyl bromoacetate (0.01068 mol) was added at 0°C. The mixture was stirred at room temperature for 1 hour, hydrolyzed with water and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:
20 cyclohexane/EtOAc 60/40; 15-40 μm). The desired fractions (F1-F4) were collected and the solvent was evaporated. Yield: 0.11g F1; 0.13g F2; 0.75g F3 and 0.8g F4. F3 was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: compound (44); mp. 152°C.
F4 was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield:
25 compound (45); mp. 147°C.



cis (compound 46)



(trans) (compound 47)

Bromomethylbenzene (0.007 mol) was added dropwise at 0°C under N₂ flow to a solution of compound (8) and compound (9) (0.0064 mol) and NaH 60% (0.007 mol) in DMF (40ml). The mixture was stirred at room temperature for 1 hour, hydrolyzed with water and extracted with EtOAc. The organic layer was separated, washed with water, dried (MgSO₄), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 70/30; 15-40 μm). The desired fractions (F1-F4) were collected and the solvent was evaporated. Yield: 0.15g F1, 0.1g F2, 0.6g F3 (23%) and 0.8g F4.

F3 was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.13g of compound (46); mp. 137°C.

F4 was crystallized from DIPE and petroleum ether. The precipitate was filtered off and dried. Yield: compound (47); mp. 130°C.

Example B7


a) 3-Chlorobenzenecarboperoxoic acid (0.088 mol) was added at 0°C to a solution of compound (48) (prepared according to example B2) (0.044 mol) in CH₂Cl₂ (200ml) and the mixture was stirred at room temperature for 12 hours. The mixture was washed with K₂CO₃ 10%. The organic layer was dried (MgSO₄), filtered off and evaporated. The residue was recrystallized from (C₂H₅)₂O. Yield : 8.2g of

cyclohexyl(3-methyl-6-quinolinyl)methanone,1-oxide (compound 49) (69%).


b) 4-Methyl benzenesulfonyl chloride (0.043 mol) was added to a solution of compound (49) (0.028 mol) in K₂CO₃ (400ml) and CH₂Cl₂ (400ml) and the mixture was stirred at room temperature for 1 hour. The mixture was extracted with CH₂Cl₂. The organic layer was dried (MgSO₄), filtered off and evaporated. The residue was recrystallized from (C₂H₅)₂O. Yield : 6.64g of 6-(cyclohexylcarbonyl)-3-methyl-2(1H)-quinolinone (compound 50) (85%); mp. 256.1°C.

Example B8

a) Preparation of




[1α(A),4α] (compound 51)



[1α(B),4α] (compound 52)

A mixture of compound (7) (0.0229 mol), hydroxylamine (0.0252 mol) and *N,N*-diethylethanamine (0.0252 mol) in ethanol (100ml) was stirred and refluxed for 6 hours, poured out into water and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue was crystallized from CH₃CN. The precipitate was filtered off and dried. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/EtOAc 80/20; 15-40μm). Two fractions were collected and the solvent was evaporated. Yielding: 2.8g of compound (51) (36%; mp. 133°C) and 3g of compound (52) (38%; mp. 142°C).

b) Preparation of




[1α(Z),4α]

(compound 53)

- Hydrazine (0.41 mol) was added at room temperature to a solution of compound (7) (0.015 mol) in ethanol (75ml). The mixture was stirred and refluxed for 1 night, poured out into water and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 98/2/0.1). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from diethyl ether. The precipitate was filtered off and dried. Yielding: 0.8g of compound (53) (15%); mp. 110°C.

Example B9

Preparation of




(compound 520)

- Procedure for compounds 400, 401, 402, 403, 404 and 405. A mixture of interm. 21 (prepared according to A11) (0.000269 mol), amantadine hydrochloride (0.000404 mol; 1.5 eq.), *N'*-(ethylcarbonimidoyl)-*N,N*-dimethyl-1,3-propanediamine

hydrochloride (0.000404 mol; 1.5 equiv.), 1-hydroxy-1*H*-benzotriazole (0.000404 mol; 1.5 equiv.) and Et₃N (0.000269 mol) in CH₂Cl₃ (2 ml) was stirred at room temperature for 12 hours. The solvent was evaporated. The residue was purified by HPLC. The product fractions were collected and the solvent was evaporated. Yield: 0.063 g of compound 520 (46.37%).

Example B10

Preparation of




(compound 233)

A mixture of intermediate 27 (0.0026 mol) and intermediate 26 (0.0026 mol) in EtOH (380 ml) and H₂SO₄ conc. (19 ml) was stirred and refluxed for 15 hours, the cooled to room temperature, poured out into ice water, basified with K₂CO₃ and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue (17.9 g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc; 80/20; 15-35μm). The pure fractions were collected and the solvent was evaporated. Yielding: 0.85 g of F1, 1.1 g F2 and 11.5 g of F3. F1 and F2 were crystallized separately from petroleum ether. The precipitate was filtered off and dried. Yielding: 0.34 g of compound 233.

Example B11

Preparation of




(compound 511)

A mixture of compound 22 (prepared according to B4) (0.004 mol) in HCl (3N) (20ml) and THF (20ml) was stirred and refluxed for 8 hours, poured out on ice, basified with NH₄OH and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue (1.2g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH; 93/7/0.5; 15-40μm). Two fractions were collected and the solvent was evaporated. Yielding: 0.5g F1 (41%) and 0.4g of F2. F1 was crystallized from petroleum ether. The precipitate was filtered off and dried. Yielding: 0.17g of compound 511 (14%).

Example B12

Preparation of




(compound 514)

A mixture of compound 524 (prepared according to B9a) (0.0018 mol) and KOH 85% (0.0094 mol) in EtOH (15ml) was stirred and refluxed for 24 hours, poured out into H₂O and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/Cyclohexane 80/20; 15-40μm). Two fractions were collected and the solvent was evaporated. Yielding: 0.35g F1 (64%) and 0.17g (SM) F1 was crystallized from diethyl ether. The precipitate was filtered off and dried. Yielding: 0.33g of compound 514 (60%) (mp.:185°C).

Example B13

Preparation of




(compound 515)

A mixture of interm. 28 (0.019 mol), 2-benzofuranylboronic acid (0.028 mol), Pd(PPh₃)₄ (0.001 mol) and BHT (a few quantity) in dioxane (25ml) and Na₂CO₃ [2] (25ml) was stirred and refluxed for 8 hours and extracted with EtOAc. The aqueous layer was basified with NH₄OH and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue (3.6g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 99/1; 15-40μm). The pure fractions were collected and the solvent was evaporated. Yielding: 1.8g (33%). This fraction was crystallized from 2-propanone/diethyl ether. The precipitate was filtered off and dried. Yielding: 0.39g of compound 515 (7%) (mp.:134°C).

Example B14

Preparation of




(compound 526)

Triethylsilane (0.0012 mol) was added slowly at room temperature to a solution of interm. 32 (0.004 mol) in CF₃COOH (5ml) and AcOH (10ml). NaBH₄ (0.0012 mol) was added portionwise under N₂ flow. The mixture was stirred at room temperature for 8 hours, poured out on ice, basified with K₂CO₃ and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue (1.2g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 99/1; 15-40μm). Two fractions were collected and the solvent was evaporated. Yielding: 0.5g F1 (43%) and 0.4g F2. F1 was dissolved in iPrOH.

HCl/iPrOH (1 eq) were added. The precipitate was filtered off and dried; Yielding: 0.32g of compound 526 (mp.: 248°C).

Example B15

Preparation of




(compound 471)

5 A mixture of interm. 33 (0.082 mol) and 3-chloro-2-ethyl-2-butenal (0.098 mol) in AcOH (200ml) was stirred and refluxed for 8 hours. The solvent was evaporated till dryness. The residue was dissolved in CH₂Cl₂ and washed with K₂CO₃ 10%. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue (27g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/EtOAc 95/5 to 92/8; 15-35μm). Two fractions were collected and the solvent
10 was evaporated. Yielding: 0.7g of F1 and 5.3g F2. F1 was crystallized from 2-propanone/diethyl ether. The precipitate was filtered off and dried. Yielding: 0.25g of compound 471 (2%) (mp.: 140°C).

Example B16

Preparation of




(compound 498)

15 *n*BuLi (0.0417 mol) was added dropwise at -78°C to a solution of interm. 35 (prepared according to A17.b) (0.0379 mol) in THF (200ml) under N₂ flow. The mixture was stirred for 30 minutes. A solution of 4-bromo-*N*-methoxy-*N*-methylbenzeneacetamide (0.0568 mol) in THF (100ml) was added dropwise at -78°C. The mixture was stirred from -78°C to 0°C, poured out into H₂O and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated till dryness. The
20 residue (20.9g) was purified by column chromatography over silica gel (eluent: toluene/EtOAc 60/40 to 50/50; 15-35μm). Two fractions were collected and the solvent was evaporated. Yielding: 4g of fraction 1 and 4g of fraction 2 (28%). Fraction 2 was crystallized from diethyl ether. The precipitate was filtered off and dried. Yielding: 1g compound 528 (m.p. 195°C).












25







Tables 1 to 8 list the compounds of formula (I-A) and (I-B) which were prepared according to one of the above examples.














Table 1































Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
54	B2	Cl	ethyl	H		-
3	B3a	Cl	ethyl	H		mp. 145°C
55	B3b	Cl	ethyl	H		mp. 131°C
56	B3b	Cl	ethyl	H		mp. 104°C
57	B3b	Cl	ethyl	H	phenylethyl	mp. 100°C
58	B3b	Cl	ethyl	H		mp. 126°C
59	B3b	Cl	ethyl	H		mp. 150°C
60	B3b	Cl	ethyl	H		mp. 138°C
61	B3b	OCH ₃	ethyl	H		-
62	B3b	OCH ₃	ethyl	H		mp. 130°C
63	B3b	OCH ₃	ethyl	H		mp. 116°C
64	B3b	Cl	ethyl	H	-(CH ₂) ₂ -O-CH ₃	mp. 82°C
65	B3b	OCH ₃	ethyl	H	1-methylcyclohexyl	mp. 82°C
66	B3b	OCH ₃	ethyl	H	3-methoxycyclohexyl	trans; mp. 94°C
67	B3b	OCH ₃	ethyl	H	3-methoxycyclohexyl	cis; mp. 108°C















Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
68	B3b	OCH ₃	ethyl	H	4-(methylethoxy)-cyclohexyl	(A), mp. 82°C
69	B3b	OCH ₃	ethyl	H	4-[C(CH ₃) ₃]cyclohexyl	cis; mp. 92°C
70	B3b	OCH ₃	ethyl	H	4-[C(CH ₃) ₃]cyclohexyl	trans; mp. 108°C
71	B3b	OCH ₃	ethyl	H	4-methylcyclohexyl	(B), mp. 92°C
72	B3b	OCH ₃	ethyl	H	4-methylcyclohexyl	(A), mp. 80°C
2	B2	Cl	ethyl	H	CH ₂ -CH(CH ₃) ₂	mp. 82°C
73	B3b	Cl	ethyl	H	-CH ₂ -O-C ₂ H ₅	mp. 82°C
48	B2	H	methyl	H	cyclohexyl	-
74	B4	I	ethyl	H		-
75	B4	I	ethyl	H		mp. 124°C
76	B4	I	ethyl	H		mp. 138°C
77	B4	I	ethyl	H		mp. 120°C
78	B4	CN	ethyl	H		mp. 128°C
79	B4	CN	ethyl	H		mp. 136°C
80	B4	CN	ethyl	H		mp. 120°C
81	B4	CN	ethyl	H		mp. 139°C
82	B4	methyl	ethyl	H		mp. 106°C
83	B4	methyl	ethyl	H		mp. 149°C
84	B4	methyl	ethyl	H		mp. 118°C














Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
85	B4	methyl	ethyl	H		mp. 180°C
86	B4	methyl	ethyl	H	phenylethyl	mp. 53°C
87	B4	methyl	ethyl	H		mp. 87°C
88	B4	methyl	ethyl	H	-CH ₂ -CH(CH ₃) ₂	mp. 68°C
89	B4	methyl	ethyl	H		mp. 120°C
31	B4	3-thiazolyl	ethyl	H	4-methoxycyclohexyl	cis; 113°C
90	B3b	OCH ₃	H	H	4-methoxycyclohexyl	trans, mp. 126°C
91	B3b	OCH ₃	H	H	4-methoxycyclohexyl	cis, mp. 100°C
92	B3b	OCH ₃	H	CH ₃	4-methoxycyclohexyl	cis; mp. 120°C
93	B3b	OCH ₃	H	CH ₃	4-methoxycyclohexyl	trans; mp. 111°C
94	B3b	OCH ₃	methyl	H	4-methoxycyclohexyl	cis, mp. 96°C
95	B3b	OCH ₃	phenyl	H	4-methoxycyclohexyl	cis; HCl (1:1), mp. 138°C
96	B3b	OCH ₃	propyl	H	4-methoxycyclohexyl	trans; mp. 118°C
97	B3b	OCH ₃	propyl	H	4-methoxycyclohexyl	cis; mp. 108°C
98	B3b	OCH ₃	methyl	H	4-methoxycyclohexyl	cis; mp. 104°C
99	B4	N(CH ₃) ₂	ethyl	H		(B); mp. 102°C
100	B3b	Cl	ethyl	H		mp. 114°C
101	B4	methyl	ethyl	H	4-butoxycyclohexyl	cis; mp. 86°C
102	B3b	Cl	ethyl	H		mp. 78°C














Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
103	B3b	Cl	ethyl	H		mp. 91°C
104	B4	N(CH ₃) ₂	ethyl	H		mp. 103°C
105	B4	N(CH ₃) ₂	ethyl	H		mp. 170°C
106	B3b	Cl	ethyl	H		mp. 137°C
107	B3b	Cl	ethyl	H		mp. 137°C
108	B4	methyl	ethyl	ethyl	4-methoxycyclohexyl	cis; mp. 91°C
109	B4	methyl	ethyl	H	4-ethoxycyclohexyl	trans; mp. 150°C
110	B4	methyl	ethyl	H		mp. 90°C
111	B4	methyl	ethyl	H		mp. 94°C
112	B4	methyl	ethyl	H		mp. 176°C
113	B4	methyl	ethyl	H		mp. 106°C
114	B4	propyl	H	H	4-methoxycyclohexyl	cis; mp. 74°C
115	B4	methyl	ethyl	H	4-ethoxycyclohexyl	cis; mp. 108°C
116	B4	methyl	ethyl	H		mp. 110°C
117	B3b	Cl	ethyl	H		mp. 124°C
118	B3b	Cl	ethyl	H		mp. 107°C
119	B3b	Cl	ethyl	H		mp. 129°C














Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
120	B4	methyl	ethyl	H		mp. 106°C
41	B3b	Cl	ethyl	H		trans; mp. 157°C
182	B3b	methyl	ethyl	H		cis; mp. 170°C
183	B3b	methyl	ethyl	H		trans; mp. 144°C
184	B3b	methyl	ethyl	H		mp. 138°C
185	B3b	Cl	ethyl	H		mp. 120°C
186	B3b	Cl	ethyl	H		
187	B3b	methyl	ethyl	H		mp. 162°C
216	B4	CC≡N	ethyl	H		mp.:160°C
217	B4	methyl	ethyl	H		ethanedioate (1:1); mp.:143°C
218	B4	I	ethyl	H		mp.:102°C
219	B4	CC≡N	ethyl	H		mp.:115°C
220	B4	Cl	ethyl	H		(A); mp.:107°C
221	B4	Cl	ethyl	H		(B); mp.:113°C

















Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
222	B4	I	ethyl	H		mp.:206°C
223	B4	Cl	ethyl	H		(trans); mp.:117°C
224	B4	methyl	ethyl	H		(A); mp.:103°C
225	B2	Cl	ethyl	H		mp.:94°C
226	B3b	Cl	ethyl	H		(trans); mp.:157°C
227	B3c	methoxy		H		mp.:204°C
228	B4	Cl	ethyl	H		(A); mp.:136°C
229	B3b	n-propyl	H	H		(trans);.HCl (1:1); mp.:150°C
230	B3b	Cl	ethyl	H		mp.:116°C
231	B3b	Cl	ethyl	H		mp.:120°C
232	B3b	Cl	ethyl	H		mp.:112°C
233	B10	<i>i</i> -propyl	H	C(=O)O-C ₂ H ₅		(cis); mp.:91°C
234	B4	methyl	ethyl	H		mp.:122°C

















Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
235	B4	methyl	ethyl	H		mp.:106°C
236	B4	methyl	ethyl	H		mp.:104°C
237	B4	methyl	ethyl	H		mp.:90°C
238	B4	methyl	H	H		(cis); mp.:80°C
239	B3b	Cl	ethyl	H		(trans); mp.:126°C
240	B3b	Cl	ethyl	H		(cis); mp.:128°C
241	B4	methyl	ethyl	H		(A); mp.:90°C
242	B4	methyl	ethyl	H		(B); mp.:110°C
243	B3b	Cl	ethyl	H		mp.:134°C
244	B3b	Cl	ethyl	H		mp.:127°C
245	B4	NHC(=O)NH ₂	ethyl	H		(cis); mp.:176°C
246	B4	methyl	ethyl	H		(B)
247	B3b	Cl	ethyl	H		mp.:92°C
248	B4	methyl	ethyl	H		(A); mp.:80°C

Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
249	B3b	Cl	ethyl	H		(B); mp.:138°C
250	B4	methyl	ethyl	H	 O- <i>n</i> propyl	(trans); mp.:118°C
251	B4	methyl	ethyl	H		(B);.HCl(1:1)
252	B3b	Cl	ethyl	H	 <i>i</i> propyl	(A)
253	B3b	Cl	ethyl	H	 <i>i</i> propyl	(B)
254	B3b	methyl	ethyl	H		mp.:74°C
255	B4	methyl	ethyl	H	 O- <i>n</i> propyl	(cis); mp.:68°C
256	B4	methyl	ethyl	H		mp.:210°C
257	B4	methyl	ethyl	H		mp.:113°C
258	B4	methyl	ethyl	H		mp.:92°C
259	B3b	methyl	ethyl	H		mp.:115°C
260	B3b	methyl	ethyl	H		mp.:60°C
261	B3b	Cl	ethyl	H		(A); mp.:86°C

Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
262	B3b	Cl	ethyl	H		(B); mp.:101°C
263	B3b	methyl	ethyl	H		mp.:130°C
264	B3b	Cl	ethyl	H		(A); mp.:124°C
265	B3b	Cl	ethyl	H		(B); mp.:126°C
266	B4	N(CH ₃) ₂	ethyl	H		(trans); mp.:102°C
267	B4	N(CH ₃) ₂	ethyl	H		(cis);.HCl(1:1); mp.:170°C
268	B4	methyl	ethyl	H		(A);.HCl(1:1); mp.:206°C
269	B4	methyl	ethyl	H		mp.:104°C
270	B3b	methyl	ethyl	H		mp.:117°C
271	B4	NHC ₂ H ₅ OCH ₃	ethyl	H		-
272	B4	methyl	ethyl	H		-
273	B4	NH ₂	ethyl	H		-
274	B3b	Cl	ethyl	H		-

Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
275	B3b	Cl	ethyl	H		mp.:99°C
276	B3b	Cl	ethyl	H		mp.:95°C
277	B4	methyl	ethyl	H		mp.:105°C
278	B3b	Cl	ethyl	H		mp.:141°C
279	B4	Cl	ethyl	H		mp.:168°C
280	B4	Cl	ethyl	H		-
281	B4	Cl	ethyl	H		mp.:140°C
282	B4	Cl	ethyl	H		mp.:169°C
283	B4	methyl	ethyl	H		mp.:96°C
284	B3b	Cl	CH ₂ N(CH ₃) ₂	H		mp.:115°C
285	B4	methyl	ethyl	H		mp.:133°C
286	B4	methyl	CH ₂ OCH ₃	H		(trans); mp.:106°C
287	B4	methyl	CH ₂ N(CH ₃) ₂	H		(cis); mp.:110°C

Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
288	B3b	Cl	<i>n</i> -propyl	H		mp.:110°C
289	B4	NH ₂	ethyl	H		mp.:218°C
290	B4	methyl	<i>n</i> -propyl	H		mp.:90°C
291	B3b	Cl	<i>n</i> -propyl	H		(cis); mp.:128°C
292	B3b	Cl	<i>n</i> -propyl	H		(trans); mp.:104°C
293	B3b	Cl	ethyl	H		mp.:106°C
294	B4	methyl	<i>n</i> -propyl	H		(cis); mp.:94°C
295	B4	methyl	CH ₂ N(CH ₃) ₂	H		mp.:83°C
296	B3b	Cl	ethyl	H		mp.:99°C
297	B3b	Cl	ethyl	H		mp.:110°C
298	B4	methyl	ethyl	H		mp.:93°C
299	B4	methyl	ethyl	H		mp.:105°C
300	B4	methyl	ethyl	H		mp.:114°C
301	B3b	methyl	ethyl	H		mp.:143°C
302	B4	methoxy	ethyl	H		mp.:93°C
303	B4	methyl	ethyl	H		mp.:82°C

Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
304	B4	n-butyl	ethyl	H		-
305	B3b	Cl	n-propyl	H		mp.:125°C
306	B1	methyl	C(=O)OC ₂ H ₅	H		mp.:136°C
307	B4	methyl	n-propyl	H		mp.:81°C
308	B4	methoxy	n-propyl	H		mp.:80°C
309	B4	I	n-propyl	H		mp.:120°C
310	B3d	methyl	ethyl	H		.HCl(1:1); mp.:129°C
311	B3b	Cl	H	H		mp.:160°C
312	B3b	Cl	H	H		(trans); mp.:145°C
313	B3b	Cl	H	H		mp.:103°C
314	B4	n-propyl	n-propyl	H		.HCl(1:1); mp.:150°C
315	B4	n-propyl	ethyl	H		.HCl(1:1)
316	B4	n-propyl	H	H		.HCl(1:1); mp.:140°C
317	B3b	Cl	H	H		mp.:168°C
318	B4	methyl	n-propyl	H		.HCl(1:1); mp.:200°C
509	B3b	Cl	ethyl	H		-



















Co. no.	Ex. no.	R ²	R ³	R ⁴	R ¹	physical data
510	B4	methyl	ethyl	H		H ₂ O(1:1)
513	B4	methyl	ethyl	H		-
516	B4	Cl	ethyl	H		mp.: 120°C
517	B4	I	ethyl	H	CH ₂ CH(CH ₃) ₂	-
518	B4	Cl	ethyl	H		-
519	B4	Cl	ethyl	H		(A+B)
521	B4	I	ethyl	H		-
522	B4	methyl	ethyl	H		(A)
1	B4	methyl	ethyl	H		(A)
525	B4	Cl	ethyl	H		
527	B4	F	ethyl	H		mp.: 116°C














Table 2





























Co. no.	Ex. no.	R ²	X	physical data
5	B3b	Cl	O	trans; mp. 120°C
121	B3b	1-piperidinyl	O	cis; HCl (1:1)
122	B3b	1-piperidinyl	O	trans; HCl (1:1); mp. 128°C

Co. no.	Ex. no.	R ²	X	physical data
123	B3b	4-thiomorpholinyl	O	cis; mp. 105°C
124	B3b	4-thiomorpholinyl	O	trans; mp. 115°C
125	B3b	4-morpholinyl	O	trans; mp. 118°C
126	B3b	4-morpholinyl	O	cis; mp. 118°C
127	B3b	-N(CH ₃) ₂	O	trans; mp. 96°C
128	B3b	-N(CH ₃) ₂	O	cis; mp. 114°C
4	B3b	Cl	O	cis; mp. 123°C
8	B3c	OCH ₃	O	trans, mp. 68°C
7	B3c	OCH ₃	O	cis, mp. 116°C
6	B4	acetyl	O	trans; mp. 108°C
129	B4	acetyl	O	cis; mp. 106°C
11	B4	NH-(CH ₂) ₂ -OCH ₃	O	trans; mp. 107°C
10	B4	NH-(CH ₂) ₂ -OCH ₃	O	cis; mp. 115°C
12	B4	NH-(CH ₂) ₂ -SCH ₃	O	cis; mp. 120°C
13	B4	NH-(CH ₂) ₂ -SCH ₃	O	trans; mp. 125°C
14	B4	-C≡C-Si(CH ₃) ₃	O	cis; mp. 114°C
16	B4	-C≡C-Si(CH ₃) ₃	O	trans; mp. 108°C
15	B4	-C≡CH	O	cis; mp. 132-133°C
17	B4	-C≡CH	O	trans; mp. 128°C
18	B4	-C≡C-CH ₂ OH	O	cis; mp. 113°C
130	B4	-C≡C-CH ₂ OH	O	trans; mp. 108°C
19	B4	F	O	cis; mp. 92-99°C
20	B4	F	O	trans; mp. 114°C
21	B4	I	O	cis; mp. 110°C
22	B4	CN	O	cis; mp. 137-138°C
26	B4	H	O	trans
23	B4	-C(=O)-OCH ₃	O	cis; mp. 91°C
24	B4	-C(=O)-OCH ₃	O	trans; mp. 99°C
25	B4	H	O	cis; mp. 88°C
27	B4	methyl	O	cis; mp. 110-112°C
131	B4	methyl	O	trans; mp. 25°C
28	B4	ethenyl	O	cis; mp. 108°C
132	B4	ethenyl	O	trans; mp. 103°C
29	B4	phenyl	O	trans; mp. 112°C

Co. no.	Ex. no.	R ²	X	physical data
30	B4	2-thienyl	O	cis; 142°C
133	B4	2-thiazolyl	O	cis; 108°C
134	B4	2-furanyl	O	cis; mp. 105°C
51	B8a	OCH ₃	N-OH	[1α(A),4α]; mp. 133°C
52	B8a	OCH ₃	N-OH	[1α(B),4α]; mp. 142°C
53	B8b	OCH ₃	NNH ₂	[1α(Z),4α]; mp. 110°C
135	B4	NH ₂	O	cis; mp. 203°C
136	B4	NH ₂	O	trans; mp. 202°C
137	B4	-C(=O)-OCH(CH ₃) ₂	O	cis; mp. 105°C
138	B4	-C(=O)-OCH(CH ₃) ₂	O	trans; mp. 88°C
38	B4	SCH ₃	O	cis; mp. 124°C
39	B4	SCH ₃	O	trans; mp. 116°C
32	B4		O	cis; mp. 130°C
139	B4	ethyl	O	cis; mp. 180°C
188	B4	NH ₂	O	cis + trans
189	B4		O	cis; mp. 154°C
190	B4		O	trans; mp. 156°C
191	B4		O	cis; mp. >260°C
192	B4		O	.H ₂ O (1:1); trans; mp. 248°C
193	B4		O	cis; mp. 224°C
194	B4		O	trans; mp. 234°C

Co. no.	Ex. no.	R ²	X	physical data
195	B4		O	cis; mp. 108°C
196	B4		O	trans; mp. 127°C
197	B4		O	cis; mp. 150°C
198	B4		O	trans; mp. 90°C
199	B4		O	LC/MS [M+H] ⁺ ; 475.4
200	B4		O	LC/MS [M+H] ⁺ ; 464.3
201	B4		O	LC/MS [M+H] ⁺ ; 523.3
202	B4		O	LC/MS [M+H] ⁺ ; 465.3
203	B4		O	LC/MS [M+H] ⁺ ; 475.4
204	B4		O	LC/MS [M+H] ⁺ ; 465.3
205	B4		O	-
319	B4		O	(cis); ethanedioate(1:1); mp.: 160°C
320	B4		O	(cis); mp.: 150°C
321	B4	methoxy	CH ₂	(cis); .HCl(1:1); mp.: 118°C
322	B4	n-butyl	O	(cis); .HCl(1:1); mp.: 158°C

Co. no.	Ex. no.	R ²	X	physical data
323	B4		O	-
324	B4		O	-
325	B4		O	-
326	B4		O	-
327	B4		O	-
328	B4		O	-
329	B4		O	-
330	B4		O	-
331	B4		O	-
332	B4		O	-
333	B4		O	-
334	B4		O	-
335	B4		O	-

Co. no.	Ex. no.	R ²	X	physical data
336	B4		O	-
337	B4		O	-
338	B4		O	-
339	B4		O	-
340	B4		O	-
341	B4		O	-
342	B4		O	-
343	B4		O	-
344	B4		O	-
345	B4		O	-
346	B4		O	-
347	B4		O	-
348	B4	CH ₂ OC(=O)CH ₃	O	(cis); mp.:74°C
349	B4		O	-

Co. no.	Ex. no.	R ²	X	physical data
350	B4		O	-
351	B4		O	-
352	B4		O	-
353	B4		O	(A);.HCl(1:2).H2O(1:1); mp.:166°C
354	B4		O	(cis)
355	B4		O	-
356	B4		O	-
357	B4		O	-
358	B4		O	-
359	B4		O	-
360	B4		O	-
361	B4		O	-
362	B4		O	-
363	B4		O	-

Co. no.	Ex. no.	R ²	X	physical data
364	B4		O	-
365	B4		O	-
366	B4		O	-
367	B4		O	-
368	B4		O	-
369	B4		O	-
370	B4		O	-
371	B4		O	-
372	B4		O	-
373	B4		O	-
374	B4		O	-
375	B4		O	-














Co. no.	Ex. no.	R ²	X	physical data
376	B4		O	-
377	B4		O	-
378	B4		O	-
379	B4		O	-
380	B4		O	-
381	B4		O	-
382	B4		O	-
383	B4		O	(cis); mp.:148°C
384	B4		O	(trans); mp.:141°C
385	B4		O	mp.:130°C
386	B4		O	(cis); mp.:140°C
387	B4		O	(trans); mp.:155°C

Table 3



Co. no.	Ex. no.	Y.	R ¹	physical data
140	B4	O		mp. 220°C
141	B4	O		mp. 213°C
142	B4	O		mp. 148°C
143	B4	O	1-methylcyclohexyl	mp. 195-210°C
144	B4	O	3-methoxycyclohexyl	cis; mp. 156°C
145	B4	O	3-methoxycyclohexyl	trans; mp. 156-163°C
146	B4	O	4-(dimethylethyl)cyclohexyl	mp. 230°C
147	B4	O	4-(methylethoxy)cyclohexyl	mp. 186°C
148	B4	O	4-methylcyclohexyl	trans; mp. 214°C
36	B4	S	4-methoxycyclohexyl	cis; mp. 224°C
37	B4	S	4-methoxycyclohexyl	trans; mp. 220°C
149	B4	O		mp. 188°C
40	B4	O		mp. 192°C
150	B4	O		cis; mp. 226°C
151	B4	O		trans; mp. 226°C
152	B4	O		mp. 213°C
153	B4	O		mp. 200°C














Co. no.	Ex. no.	Y.	R ¹	physical data
154	B4	O		mp. 210°C
155	B4	O	4,4-dimethylcyclohexyl	mp. 242°C
388	B4	O	CH ₂ CH(CH ₃) ₂	mp. 189°C
389	B4	O		mp. 228°C
390	B4	O		mp. 197°C
391	B4	O		mp. 145°C
392	B4	O		mp. 192°C
393	B4	O		(B); mp.:224°C
394	B4	O		(A); mp.:201°C
395	B4	O		(A); mp.:207°C
396	B4	O		mp.:212°C
397	B4	O		(B); mp.:238°C
398	B4	O		mp.:234°C
399	B4	O		(cis); mp.:192°C

Table 4



Co. no.	Ex. o.	R ³	R ⁴	R ⁵	R	physical data
156	B4	ethyl	H	H	OCH ₃	trans; mp. 252°C
157	B4	H	H	H	OCH ₃	(cis + trans); mp. 244°C
158	B4	H	methyl	H	OCH ₃	cis; mp. >260°C
159	B4	methyl	H	H	OCH ₃	cis; mp. 254°C
160	B4	methyl	H	H	OCH ₃	trans; mp. >260°C
161	B4	propyl	H	H	OCH ₃	mp. 208°C
162	B4	propyl	H	H	OCH ₃	trans; mp. 232°C
9	B4	ethyl	H	H	OCH ₃	cis; mp. 224-226°C
43	B5	ethyl	H	CH ₃	OCH ₃	trans; mp. 116°C
42	B5	ethyl	H	CH ₃	OCH ₃	cis; mp. 125°C
44	B6	ethyl	H	CH ₂ -COOC ₂ H ₅	OCH ₃	cis; mp. 152°C
45	B4	ethyl	H	CH ₂ -COOC ₂ H ₅	OCH ₃	trans; mp. 147°C
46	B4	ethyl	H	benzyl	OCH ₃	cis; mp. 137°C
47	B4	ethyl	H	benzyl	OCH ₃	trans; mp. 130°C
50	B7	methyl	H	H	H	mp. 256.1°C
163	B4	ethyl	ethyl	H	OCH ₃	cis; mp. 221°C
164	B4	ethyl	ethyl	H	OCH ₃	cis; mp. 221°C
165	B4	ethyl	ethyl	H	OCH ₃	trans; mp. 215°C
166	B4	ethyl	H		OCH ₃	LC/MS [M+H] ⁺ ; 429.4
167	B4	ethyl	H		OCH ₃	LC/MS [M+H] ⁺ ; 451.3
168	B4	H	H	H	OCH ₃	cis; mp. 106°C
169	B4	ethyl	H		OCH ₃	LC/MS [M+H] ⁺ ; 409.3















Co. no.	Ex. o.	R ³	R ⁴	R ⁵	R	physical data
400	B9	ethyl	H		OCH ₃	-
401	B9	ethyl	H		OCH ₃	-
402	B9	ethyl	H		OCH ₃	-
403	B9	ethyl	H		OCH ₃	-
404	B9	ethyl	H		OCH ₃	-
405	B9	ethyl	H		OCH ₃	-
406	B4	ethyl	H		OCH ₃	-
407	B4	ethyl	H		OCH ₃	-
408	B4	ethyl	H		OCH ₃	-
409	B3b		H	H	OCH ₃	mp.:168°C
410	B4	CH ₂ OCH ₃	H	H	OCH ₃	mp.:194°C
508	B4	ethyl	H		OCH ₃	-
520	B9	ethyl	H		OCH ₃	-

Table 5



























Co. no.	Ex. no.	R ⁴	R ¹	X	physical data
33	B4	H	methoxycyclohexyl	CH	cis; mp. 224°C
34	B4	H	methoxycyclohexyl	CH	trans; mp. 185°C
35	B4	H	methoxycyclohexyl	N	cis; mp. 160-172°C
170	B4	H	methoxycyclohexyl	N	trans; mp. 146°C
171	B4	H		N	(B); mp. 165°C
172	B4	H	methylecyclohexyl	N	cis+trans; mp. 143°C
173	B4	ethyl	methoxycyclohexyl	N	cis; mp.:126°C
411	B4	H		N	mp.:109°C
412	B4	H		N	mp.:180°C
413	B4	H		N	(A)
414	B4	H		N	mp.:156°C



Table 6




Co. no.	Ex. no.	R	L	physical data
49	B7	H		-
174	B3b	OCH ₃		cis; mp.115°C
175	B3b	OCH ₃		trans; mp.141°C
176	B3b	OCH ₃		cis; mp.149°C
177	B3b	OCH ₃		mp.126°C
178	B3b	OCH ₃		trans; mp.160°C
179	B3b	OCH ₃		cis; mp.119°C
180	B3b	OCH ₃		trans; mp.124°C
181	B3b	OCH ₃		trans; mp.92°C
206	B3b	OCH ₃		cis; m.p. 144°C



















Co. no.	Ex. no.	R	L	physical data
207	B3b	OCH ₃		trans; m.p. 125°C
208	B3b	OCH ₃		cis; m.p. 127°C
209	B3b	OCH ₃		cis; m.p. 101°C
210	B3b	OCH ₃		cis; m.p. 104°C
211	B3b	OCH ₃		trans; m.p. 134°C
212	B4	OCH ₃		cis; m.p. 141°C
213	B4	OCH ₃		trans; m.p. 215°C
214	B4	OCH ₃		cis; m.p. 139°C
215	B3b	OCH ₃		trans
415	B3b	OCH ₃		(cis); mp.:136°C
416	B3b	OCH ₃		(cis)





























Co. no.	Ex. no.	R	L	physical data
417	B4	OCH ₃		(cis); mp.:149°C
418	B3b	OCH ₃		(trans); mp.:132°C
419	B4	OCH ₃		(cis); mp.:217°C
420	B3b	OCH ₃		(cis); .HCl(1:1); mp.:200°C
421	B4	OH		(cis); mp.:215°C
422	B4	OH		(trans); mp.:178°C
423	B3b	OCH ₃		mp.:160°C
424	B3b	OCH ₃		(cis); mp.:106°C
425	B3b	OCH ₃		(trans); mp.:120°C
426	B3b	OCH ₃		(cis); mp.:121°C
427	B3b	H		mp.:156°C
428	B3b	OCH ₃		(cis); mp.:156°C
429	B3b	OCH ₃		(trans); mp.:197°C






























Co. no.	Ex. no.	R	L	physical data
430	B3b	CH ₃		(B)
431	B3b	CH ₃		(A)






























5 Table 7































Co. no.	Ex. no.	R ¹	L	physical data
432	B16			mp.:128°C
433	B4			mp.:175°C
434	B4			mp.:170°C
435	B4			mp.:103°C
436	B4			mp.:151°C
437	B4			(trans); mp.:110°C
438	B4			mp.:150°C
439	B4			mp.:150°C
440	B4			(cis)

Co. no.	Ex. no.	R ¹	L	physical data
441	B4			mp.:166°C
442	B4			mp.:173°C
443	B4			mp.:208°C
444	B4			mp.:149°C
445	B4			mp.:133°C
446	B3b			mp.:150°C
447	B3b			mp.:165°C
448	B3b			mp.:147°C
449	B3b			mp.:154°C
450	B3b			mp.:157°C
451	B4			mp.:190°C
452	B4			mp.:187°C
453	B3b			mp.:200°C
454	B3b			mp.:160°C

Co. no.	Ex. no.	R ¹	L	physical data
455	B3b			mp.:139°C
456	B3b			(A); mp.:174°C
457	B3b			(B); mp.:160°C
458	B3b			mp.:184°C
459	B4			-
460	B4			mp.:134°C
461	B4			(B); mp.:156°C
462	B4			mp.:153°C
463	B3b			mp.:161°C
464	B4			mp.:135°C
465	B4			mp.:131°C
466	B3b			.HCl(1:1); mp.:206°C
467	B3d			mp.:142°C
468	B4			.hydrate(1:1); mp.:104°C
469	B3b	dimethylethyl		mp.:104°C

Co. no.	Ex. no.	R ¹	L	physical data
470	B3b			mp.:161°C
472	B3b			mp.:144°C
473	B4			mp.:143°C
474	B4			mp.:196°C
475	B4			mp.:162°C
476	B4			mp.:171°C
477	B4			mp.:155°C
478	B2	trimethylmethyl		mp.:124°C
479	B4			(A); mp.:146°C
480	B4			(B); mp.:162°C
481	B4			(A); mp.:129°C
482	B4			mp.:115°C
483	B2			mp.:187°C
484	B2			mp.:162°C
485	B4			(A); mp.:130°C

Co. no.	Ex. no.	R ¹	L	physical data
486	B4			(A); mp.:124°C
487	B4			(B); mp.:128°C
488	B4			mp.:85°C
489	B2			mp.:150°C
490	B4			(A); mp.:117°C
491	B2			mp.:220°C
492	B4			mp.:136°C
493	B2			mp.:131°C
494	B4			(A); mp.:125°C
495	B4			mp.:135°C
496	B4			mp.:139°C
497	B4			mp.:127°C
498	B16			mp.:195°C
499	B2			mp.:201°C



























Co. no.	Ex. no.	R ¹	L	physical data
500	B3b			mp.:143°C
501	B3b			mp.:137°C
502	B2			mp.:210°C
503	B3d			mp.:134°C
504	B2			mp.:163°C
505	B4			mp.:142°C
506	B2			mp.:139°C
507	B4			mp.:171°C
512	B3b			-
523	B3b			-

Table 8:


Co. no.	Ex. no.	Structure	physical data
511	B11		-

Co. no.	Ex. no.	Structure	physical data
514	B12		-
515	B13		-
524	B9a		mp.:185°C
471	B15		(E)
526	B14		.HCl(1:1)

C. Preparation of radioactively labelled compounds

C.1 [³H]-labelled compounds

5



- 10 To a carefully measured amount of palladium on carbon (10 %, 0.872 mg) was added a solution of compound 498 (I, 0.919 mg, 2.4 μmol) and triethylamine (0.92 μl, 6.6 μmol) in sodium-dried tetrahydrofuran (175 μl). The reaction flask was connected to a tritiation manifold system and the reaction mixture was carefully degassed. Tritium gas (19.5 Ci at a pressure of 1017 mbar) was generated from uranium tritide and was

allowed onto the at room temperature stirred reaction mixture. After 30 min, the reaction mixture was frozen with liquid nitrogen and the excess of tritium gas was recaptured onto uranium sponge. The solvent was lyophilized from the reaction mixture. Methanol (100 μ l) was introduced and lyophilized in order to remove labile tritium. This procedure was repeated twice more. The residue was taken up in ethanol, filtered over a GHP Acrodisk 13 mm syringe filter and depleted with ethanol to a total volume of 50.0 ml. It contained 71 mCi of radioactivity with [3 H]-compound 528 (II) at a 67 % radiochemical purity. From this amount, a fraction (5.0 ml) was taken and thoroughly purified in portions via preparative HPLC (Kromasil KR 100-10, column dimensions 4.6 mm ID x 300 mm). UV detection took place at 265 nm. Elution was performed isocratically with water-methanol-acetonitrile-diisopropylamine (47:26.5:26.5:0.2; v/v/v/v) at a flow rate of 2.0 ml/min. The product containing fractions were combined and concentrated under vacuum at 30°C. The residue was dissolved in ethanol (5.0 ml) and concentrated again. This procedure was repeated twice more. The remaining residue was finally dissolved in ethanol (20.0 ml) and stored as such. The batch contained [3 H]-compound 528 (II) with a total radioactivity of 3.83 mCi at a purity > 98 % and at a specific activity of about 25 Ci/mmol.

D. Pharmacological examples

D1. Signal transduction at the cloned rat mGlu1 receptor in CHO cells

CHO cells expressing the mGlu1 receptor were plated in precoated black 96-well plates. The next day, the effect of the present compounds on glutamate-activated intracellular Ca^{2+} increase was evaluated in a fluorescent based assay. The cells were loaded with Fluo-3 AM, plates were incubated for 1 hour at room temperature in the dark, cells were washed and the present compounds were added onto the cells for 20 minutes. After this incubation time, the glutamate-induced Ca^{2+} rise was recorded for each well in function of time using the Fluorescent Image Plate Reader (FLIPR, Molecular Devices Inc.). Relative fluorescence units were recorded and average data graphs of quadruple wells were obtained. Concentration-response curves were constructed based on peak fluorescence (maximum signal between 1 and 90 secondes) for each concentration of tested compound. pIC_{50} values are the $-\log$ values of the concentration of the tested compounds resulting in 50% inhibition of the glutamate-induced intracellular Ca^{2+} rise.

The compounds according to the present invention exhibited a pIC_{50} value of at least 5. The compounds that are included in the Tables 1-8 exhibited a pIC_{50} value of at least 6.

A particular group of compounds exhibited a pIC₅₀ value between 7 and 8. It concerns the compounds listed in Table 9.

Table 9:

Com.nr.	pIC ₅₀
463	7.98
441	7.95
334	7.95
22	7.94
421	7.94
15	7.93
440	7.93
139	7.93
178	7.92
338	7.91
87	7.90
462	7.90
394	7.90
423	7.89
21	7.87
220	7.87
479	7.86
483	7.86
485	7.84
9	7.84
110	7.84
248	7.84
341	7.83
163	7.81
433	7.79
238	7.79
224	7.78
437	7.78
498	7.78
449	7.77
242	7.76

Com.nr.	pIC ₅₀
281	7.63
487	7.63
299	7.63
431	7.61
98	7.57
464	7.57
446	7.56
251	7.55
484	7.54
494	7.53
128	7.52
344	7.52
161	7.49
298	7.48
454	7.45
456	7.45
277	7.44
91	7.43
356	7.42
229	7.41
333	7.41
326	7.41
369	7.40
430	7.39
435	7.38
35	7.36
228	7.36
429	7.36
117	7.35
291	7.35
313	7.35

Com.nr.	pIC ₅₀
89	7.25
108	7.25
373	7.25
255	7.23
527	7.23
303	7.22
296	7.22
221	7.21
193	7.21
14	7.20
131	7.19
438	7.19
148	7.18
496	7.18
236	7.17
332	7.17
481	7.16
191	7.16
457	7.14
20	7.14
145	7.13
268	7.13
512	7.13
474	7.13
10	7.11
307	7.11
426	7.11
466	7.10
97	7.08
83	7.08
434	7.08

Com.nr.	pIC ₅₀
346	7.74
182	7.73
486	7.73
447	7.72
7	7.72
175	7.71
475	7.71
480	7.71
213	7.70
239	7.70
241	7.67
461	7.65
115	7.64
445	7.63



Com.nr.	pIC ₅₀
280	7.34
460	7.34
482	7.34
343	7.33
425	7.32
473	7.32
287	7.31
448	7.31
243	7.29
323	7.28
159	7.28
289	7.27
184	7.26
436	7.26







Com.nr.	pIC ₅₀
300	7.08
199	7.07
290	7.06
112	7.05
348	7.05
286	7.03
442	7.03
422	7.02
283	7.02
318	7.02
36	7.00
396	7.00







A particular group of compounds exhibited a pIC₅₀ value of at least 8. It concern the compounds listed in Table 10.







Table 10 :







5





Comp. nr.	Structure	pIC ₅₀
416	 (CIS)	8.587
27	 (CIS)	8.527

Comp. nr.	Structure	pIC50
174	 (CIS)	8.49
506		8.48
25	 (CIS)	8.45
4	 (CIS)	8.4
19	 (CIS)	8.38
429	 (CIS)	8.38

Comp. nr.	Structure	pIC50
424	 <p>(CIS)</p>	8.355
176	 <p>(CIS)</p>	8.33
210	 <p>(CIS)</p>	8.315
114	 <p>(CIS)</p>	8.28
488		8.27
504		8.27

Comp. nr.	Structure	pIC50
477		8.25
432		8.237
214	 (CIS)	8.233
465		8.145
135	 (CIS)	8.14
420	 (CIS) Hydrochloride (1:1)	8.135

Comp. nr.	Structure	pIC50
292	 (CIS)	8.13
427		8.115
208	 (CIS)	8.095
419	 (CIS)	8.065
455		8.055
418	 (TRANS)	8.045

Comp. nr.	Structure	pIC50
497		8.025
439		8.023
237		8.01
499		8

D2. In vitro binding experiments with a [³H]-radiolabelled compound according to the invention

As [³H]-radiolabelled compounds are used: compound 528, hereafter named as
5 [³H]Compound A, which is the tritium-radiolabelled equivalent of compound 432.

In the next paragraphs, a study will be disclosed illustrating the use of radiolabelled compounds according to the invention.

Materials

10 All cell culture reagents were obtained from Invitrogen (Carlsbad, USA). Glutamate was obtained from Aldrich Chemical Company (Milwaukee, WI) ; [³H]quisqualate (29 Ci/mmol), [³H]Ro 48-8587 (53 Ci/mmol), *myo*-[³H]-inositol (22 Ci/mmol) and [³⁵S]GTPγS (1030 Ci/mmol) were obtained from Amersham (Paisley, UK). [³H]MK-801 (22.5 Ci/mmol) and [³H]CGP39653 (20-50 Ci/mmol) were obtained from NEN
15 (Zaventem, Belgium). GDP was obtained from Boehringer Mannheim (Basel, Switzerland) and glycine from BioRad (CA, USA). [³H]L689560 (10-30 Ci/mmol), [³H]LY341495 (34.61 Ci/mmol), [³H]MPEP (50.2 Ci/mmol), (S)-4C3HPG, (1S,3R)-ACPD, (S)-3,5-DHPG, (S)-4CPG, AIDA, MCPG, MPEP, CPCCOEt, L-SOP and L-quisqualic acid were purchased from Tocris Cookson (Essex, UK). BAY 36-7620,
20 NPS 2390 and phencyclidine were synthesized in-house. Fluo-3-AM and pluronic acid were obtained from Molecular Probes (Leiden, The Netherlands). Probenecid, strychnine, D-serine and Triton X-100 were purchased from Sigma-Aldrich (Steinheim, Germany). All other reagents were from Merck (Darmstadt, Germany).

Cell transfection and culture

25 L929sA cells stably expressing the human mGlu1a receptor were obtained as described in Lavreysen et al., Mol. Pharmacol. **61**:1244-1254, 2002 and were cultured in Glutamax-I medium supplemented with 10% heat inactivated dialysed foetal calf serum, 0.1 mg/ml streptomycin sulphate and 100 units/ml penicillin. CHO-dhfr⁻ cells stably expressing the rat mGlu1a, -2, -3, -4, -5 and -6 receptor were a kind gift from S.
30 Nakanishi (Tokyo University, Japan) and were grown in DMEM with Glutamax-I with 10% heat inactivated dialysed foetal calf serum, 0.4 mM L-prolin, 0.2 mg/ml streptomycin sulphate and 200 units/ml penicillin. Cells were kept in an atmosphere of 37°C and 5% CO₂.

Intracellular Ca^{2+} response in rat and human mGlu1a receptor expressing cells and in rat mGlu5 receptor expressing cells.

Intracellular calcium ion levels ($[\text{Ca}^{2+}]_i$) in human mGlu1a receptor expressing L929sA cells were measured using the Fluorometric Imaging Plate Reader (FLIPR, Molecular
5 Devices, CA, USA), as described in Lavreysen et al., Mol. Pharmacol. **61**:1244-1254, 2002. The same procedure was followed for CHO-dhfr⁻ cells expressing the rat mGlu1a receptor. For the rat mGlu5 receptor, cells were seeded at 30.000 cells/well 2 days before the experiment.

IP response in rat mGlu1a receptor expressing CHO-dhfr⁻ cells

10 IP accumulation was measured as described in Lavreysen et al., Mol. Pharmacol. **61**:1244-1254, 2002. Briefly, cells were seeded at 30,000 cells/well in 24-well plates and were labelled with 2.5 $\mu\text{Ci/ml}$ *myo*- $[\text{}^3\text{H}]$ inositol overnight. On the day of the experiment, cells were washed and incubated for 10 min with 10 mM LiCl. After 30 min incubation with increasing concentrations of $[\text{}^3\text{H}]$ Compound A, 1 N HClO_4 was
15 added and plates were put at 4°C. KOH/phosphate solution and a solution containing 30 mM $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$ and 3 mM EDTA were added prior to application to ion exchange chromatography.

Membrane preparation from CHO-dhfr⁻ cells expressing the rat mGlu1a, -2, -3, -4, -5 and -6 receptor

20 Confluent cells were washed in ice-cold phosphate-buffered saline and stored at -20°C until membrane preparation. After thawing, cells were suspended in 50 mM Tris-HCl, pH 7.4 and collected through centrifugation for 10 min at 23,500 g at 4°C. The cells were lysed in 10 mM hypotonic Tris-HCl, pH 7.4. After recentrifugation for 20 min at 30,000 g at 4°C, the pellet was homogenized with an Ultra Turrax homogenizer in 50
25 mM Tris-HCl, pH 7.4. Protein concentrations were measured by the Bio-Rad protein assay using bovine serum albumin as standard.

$[\text{}^{35}\text{S}]\text{GTP}\gamma\text{S}$ binding to membranes from CHO-dhfr⁻ cells expressing the rat mGlu2, -3, -4 and -6 receptor

30 Membranes were thawed on ice and diluted in 10 mM HEPES acid, 10 mM HEPES salt, pH 7.4, containing 100 mM NaCl, 3 mM MgCl_2 , 3 μM GDP and 10 $\mu\text{g/ml}$ saponine. Assay mixtures contained 10 μg of membrane protein and were pre-incubated with compounds or buffer for 5 min at 37°C. Then, glutamate was added and the assay mixtures were further incubated for 30 min at 37°C. $[\text{}^{35}\text{S}]\text{GTP}\gamma\text{S}$ was added to a final concentration of 0.1 nM for another 30 min at 37°C. Reactions were

terminated by rapid filtration through Unifilter-96 GF/B filter plates (Packard, Meriden, CT) using a 96-well Packard filtermate harvester. Filters were washed 2 times with ice-cold 10 mM NaH₂PO₄/10 mM Na₂HPO₄ buffer, pH 7.4. Filter-bound radioactivity was counted in a Microplate Scintillation and Luminescence Counter from Packard.

5 Radioligand binding to rat mGlu1a receptor CHO-dhfr⁻ membranes

[³H]Compound A-binding. After thawing, the membranes were homogenized using an Ultra Turrax homogenizer and suspended in ice-cold binding buffer containing 50 mM Tris-HCl (pH 7.4), 1.2 mM MgCl₂, 2 mM CaCl₂, unless otherwise indicated. Ligand saturation experiments were performed at apparent binding equilibrium (30 min
10 incubation) with 20 µg membrane protein and 10 concentrations (0.1, 0.2, 0.3, 0.4, 0.5, 1, 2, 2.5, 5 and 10 nM) of radioligand. Non-specific binding was estimated in the presence of 1 µM compound 135. The incubation was stopped by rapid filtration under suction over GF/C glass-fibre filters using a manual 40-well filtration manifold. To measure association kinetics, membranes were incubated at 4°C, 25°C or 37°C in the
15 presence of 2.5 nM [³H]Compound A for 2, 5, 10, 15, 20, 30, 45, 60, 90 or 120 min, then terminated by rapid filtration using a manual 40-well filtration unit. Dissociation kinetics were measured by adding, at different times before filtration 1 µM compound 135 to membranes preincubated for 30 min at 4°C or 25°C in the presence of 2.5 nM [³H]Compound A. The filters were transferred to scintillation vials and, after the
20 addition of Ultima-Gold MV, the radioactivity collected on the filters was counted in a Packard scintillation counter. For inhibition experiments, assay mixtures were incubated for 30 min at 4°C in a volume of 0.5 ml containing 10-20 µg membrane protein, appropriate concentrations of test compounds and 2.5 nM [³H]Compound A. Non-specific binding was defined as above. Filtration was performed using Unifilter-
25 96 GF/C filter plates and a 96-well Packard filtermate harvester. After the addition of microscint-O, radioactivity on the filters was counted in a Microplate Scintillation and Luminescence Counter from Packard.

[³H]quisqualate binding. Thawed membranes were homogenized and suspended in ice-cold binding buffer. For saturation experiments, 30 µg of membrane protein was
30 incubated for 1 h at 25°C with 10 concentrations (1, 2, 5, 10, 20, 40, 60, 90, 120 and 150 nM) of [³H]quisqualate. Non-specific binding was determined in the presence of 1 mM L-glutamate. Bound and free radioligand was separated by rapid filtration over GF/C glass-fibre filters using a manual 40-well filtration manifold. For inhibition experiments, 30 µg membrane protein was incubated for 1 h at 25°C in a volume of 0.5
35 ml containing appropriate concentrations of test compounds and a final concentration

of 10 nM [^3H]quisqualate. Filtration was performed using Unifilter-96 GF/C filter plates and a Packard filtermate harvester. Radioactivity trapped on the filters was counted as above.

Radioligand binding to membranes from CHO-dhfr⁻ cells expressing the rat mGlu2, -3, -4, -5 and -6 receptor

After thawing, the membranes were homogenized using an Ultra Turrax homogenizer and suspended in ice-cold binding buffer containing 50 mM Tris-HCl (pH 7.4), 1.2 mM MgCl_2 , 2 mM CaCl_2 . For [^3H]Compound A binding, 20 to 160 μg membrane protein and a final concentration of 20 nM [^3H]Compound A was used. As indicated in the results section, different blanks were used to define non-specific binding. Incubation time and temperature as well as filtration were as described for rat mGlu1a receptor CHO-dhfr⁻ membranes. Expression of rat mGlu2, -3, -5 and mGlu6 receptors was confirmed by specific binding of [^3H]LY341495 (mGlu2, -3 and -6) or [^3H]MPEP (mGlu5). For [^3H]LY341495 binding, 1 nM (mGlu2 and mGlu3) or 10 nM (mGlu6) [^3H]LY341495 was used. Non-specific binding was determined using 1 mM glutamate. Assay mixtures were incubated for 30 min (mGlu2 and mGlu3) or 60 min (mGlu6) at 4°C. Incubation was stopped by filtration over GF/B glass fibre filters (Whatman, England) using a manual 40-well filtration manifold. For rat mGlu5 receptor CHO-dhfr⁻ membranes, 10 nM [^3H]MPEP and 10 μM MPEP, to reveal non-specific binding, were used. Incubation was performed at 4°C for 30 min. Bound and free radioligand were separated over GF/C glass-fibre filters (Whatman, England) using a 40-well filtration unit.

[^3H]Compound A binding to rat brain membranes.

Tissue preparation. Male Wistar rats (~200 g) were sacrificed by decapitation. The brains were rapidly removed and cortex, hippocampus, striatum and cerebellum were immediately dissected. The fresh tissue was homogenized with an Ultra Turrax in 20 volumes of 50 mM Tris-HCl, pH 7.4 and tissue was centrifuged at 23,500 g for 10 min. After homogenisation using a DUAL homogeniser, membranes were washed twice by centrifugation at 23,500 g for 10 min. The final pellet was suspended in 10 volumes of 50 mM Tris-HCl, pH 7.4 and frozen at -80°C.

In vitro binding assay. After thawing, membranes from rat cortex, cerebellum, striatum and hippocampus were rehomogenized using a DUAL and suspended in ice-cold binding buffer containing 50 mM Tris-HCl, 1.2 mM MgCl_2 , 2 mM CaCl_2 , pH 7.4. The binding assay was carried out in a total volume of 0.5 ml containing 2.5 nM

[³H]Compound A and a membrane aliquot corresponding to 40 µg for cerebellar membranes, 60 µg for hippocampal membranes, 80 µg for striatal membranes or 150 µg for cortical membranes. Specific binding was calculated as the difference between the total binding and the binding measured in the presence of 1 µM compound 135.

- 5 After incubation for 30 min at 4°C, the labelled membranes were washed and harvested by rapid vacuum filtration over Whatman GF/C glass-fibre filters using a 40-well filtration manifold and radioactivity collected on the filters was counted as above.

[³H]Ro 48-8587, [³H]L689560, [³H]CGP39653 and [³H]MK-801 binding to rat brain membranes.

- 10 **Tissue preparation.** Male Wistar rats (~200 g) were sacrificed by decapitation. The brains were rapidly removed and forebrain was dissected. The tissue was homogenized with an Ultra Turrax in 20 volumes of ice-cold H₂O and was centrifuged at 48,000 g for 20 min. After homogenisation using a DUAL homogeniser, membranes were washed by centrifugation at 48,000 g for 10 min. The pellet was then suspended in 20 volumes
15 of 50 mM Tris-HCl, pH 7.4 containing 0.04% Triton X-100 and again centrifuged at 48,000 g for 20 min. The final pellet was frozen at -80°C.

- In vitro binding assay.** At the day of the experiment, the pellet was thawed, washed and rehomogenised using a DUAL in ice-cold 50 mM Tris-acetate, pH 7.4. Assay conditions for the different radioligands were as follows. The final concentration of
20 membrane in the assay was 20 mg/ml (wet weight) for [³H]Ro 48-8587 and [³H]L689560 and was 10 mg/ml (wet weight) for [³H]CGP39653 and [³H]MK-801. Radioligand concentrations of 2 nM [³H]Ro 48-8587, 2 nM [³H]L689560, 2 nM [³H]CGP39653 and 3 nM [³H]MK-801 were used. Incubation was performed in the presence of 1 mM KSCN for [³H]Ro 48-8587, 100 µM strychnine for [³H]L689560 and
25 1 µM glycine + 1 µM glutamate for [³H]MK-801 binding. Non-specific binding was determined in the presence of 1 mM glutamate for [³H]Ro 48-8587 and [³H]CGP39653 binding. For [³H]L689560 and [³H]MK-801 binding, 100 µM D-serine or 10 µM phencyclidine were used to define non-specific binding, respectively. Assays were incubated for 1 h at 37°C, 2 h at 4°C, 30 min at 25°C and 1 h at 4°C for [³H]Ro 48-
30 8587, [³H]L689560, [³H]CGP39653 and [³H]MK-801 binding, respectively. After incubation, bound and free radioligand was separated using a 40-well filtration manifold. Radioactivity collected on the filters was counted as above.

[³H]Compound A-binding and autoradiography on rat brain sections.

Tissue preparation. Male Wistar rats (200 g) were sacrificed by decapitation. Brains were immediately removed from the skull and were rapidly frozen in dry-ice-cooled 2-methylbutane (-40°C). Brains were then stored at -70°C until sectioning. Twenty-micrometer-thick sagittal sections were cut using a Leica C3050 cryostat microtome (Leica Microsystems, Wetzlar, Germany) and thaw-mounted on SuperFrost Plus microscope slides (Menzle-glaser, Germany). The sections were then kept at -70 °C until use.

Receptor autoradiography. Sections were thawed and dried under a stream of cold air, preincubated (3 x 5 min) in 50 mM Tris-HCl, 1.2 mM MgCl₂, 2 mM CaCl₂, 0.1% BSA pH 7.4 at room temperature. Sections were then incubated for 60 min at room temperature, in buffer containing 50 mM Tris-HCl, 1.2 mM MgCl₂, 2 mM CaCl₂, 0.1% BSA (pH 7.4) and 1.5 nM [³H]Compound A. Non-specific binding was determined by addition of 1 μM compound 135 in the incubation buffer. After the incubation, the excess of radioligand was washed off (3 x 5 min) in ice-cold buffer containing 50 mM Tris-HCl, 1.2 mM MgCl₂ and 2 mM CaCl₂, followed by a rapid dip in cold distilled water. The sections were dried under a stream of cold air and then exposed to [³H]Hyperfilm (Amersham, UK) for 6 weeks at room temperature. The films were developed manually in Kodak D19 and fixed with Kodak Readymatic. Some sections were exposed to a Fuji Imaging Plate for 2 days at room temperature and scanned using a Fujix Bass 2000 phosphoimager.

Data analysis and statistics

Data analysis was performed using the GraphPad Prism program (GraphPad Prism Software, Inc., San Diego, CA). Saturation binding experiments were analysed using a non-linear regression analysis. Inhibition curves were fitted using non-linear regression analysis fitting the one-site competition equation: $Y = \text{Bottom} + ((\text{Top} - \text{Bottom}) / (1 + 10^{X - \text{LogIC}_{50}}))$. K_i values were calculated using the Cheng-Prusoff equation: $K_i = \text{IC}_{50} / [1 + ([C] / K_D)]$ where C is the concentration of radioligand and K_D is the dissociation constant of the radioligand (Cheng and Prusoff, Biochem. Pharmacol. **22**, 3099-3108, 1973). The observed on (k_{on}) and off (k_{off}) rate were calculated from association-dissociation curves using the one-phase-exponential association and decay equations in the Prism program, respectively. k_{on} was calculated by subtracting k_{off} from k_{ob} and dividing by the radioligand concentration. The two-tailed Student's t-test was used for statistical evaluation of the binding data: * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$.

The Dunnett's t-test following a 2-way analysis of variance (with as factors compound concentration and experiment) were used to analyse the data from the IP experiments.

Results

5 **Selectivity and mode of antagonism of Compound A for the mGlu1 receptor.** In CHO-dhfr⁻ cells expressing the rat mGlu1a receptor, compound A inhibited the glutamate-induced increase in $[Ca^{2+}]_i$ with an IC_{50} value of 21.6 ± 5.0 nM (n=4; Figure 1A) and appeared to be about 8 times more potent than the recently described specific mGlu1 receptor antagonist BAY 36-7620 ($IC_{50} = 161 \pm 38$ nM, n=3) and 500 times
10 more potent than CPCCOEt ($IC_{50} = 10.3 \pm 0.8$ μ M, n=3), tested in the same assay. For the human mGlu1a receptor, compound A had an IC_{50} value of 10.4 ± 4.7 nM (n=3). Compound A did not inhibit glutamate-induced Ca^{2+} signaling of the rat mGlu5 receptor expressed in CHO-dhfr⁻ cells, tested up to a concentration of 10 μ M. IC_{50} values for inhibition of glutamate (30 μ M)-induced $[^{35}S]GTP\gamma S$ activation were above
15 30 μ M at recombinant rat mGlu2, -3, -4 or -6 receptors. In $[^{35}S]GTP\gamma S$ assays, compound A did not exhibit agonist activity towards any of the mGlu receptors up to a concentration of 30 μ M. In addition, it was investigated whether compound A could act as a positive allosteric modulator on one of these mGlu receptor types. For this, we performed glutamate concentration-response curves by adding glutamate alone or
20 together with 10 μ M compound A. $[^{35}S]GTP\gamma S$ assays on recombinant rat mGlu2, -3, -4 or -6 receptors showed that the glutamate EC_{50} was not altered and that the glutamate E_{max} value was not increased upon addition of compound A. The EC_{50} and E_{max} value of glutamate-induced intracellular Ca^{2+} mobilization also did not change in cells expressing the rat mGlu5 receptor when compound A was added together with
25 glutamate (data not shown). Together these data exclude agonist, antagonist or positive allosteric action on mGlu2, -3, -4, -5 and -6 receptors. Radioligand binding studies on rat forebrain using $[^3H]Ro-488587$, $[^3H]L689560$, $[^3H]CGP39653$ and $[^3H]MK-801$ revealed that compound A did not bind to the AMPA receptor, nor did it bind to the glycine, glutamate or channel pore site of the NMDA receptor (tested up to a
30 concentration of 10 μ M), respectively. To analyse how compound A inhibits glutamate activation of the mGlu1a receptor, mobilization of Ca^{2+} in response to glutamate was compared in the absence and presence of compound A (Figure 1B). The presence of compound A not only caused a right-ward shift in the concentration-response curve of glutamate, but also resulted in a dramatic decrease in the maximal response evoked by
35 the agonist, revealing that antagonism by compound A was non-competitive. Complete inhibition of mGlu1a receptor-mediated signalling was observed in the presence of 100

nM-1 μ M compound A. To investigate whether compound A could act as an inverse agonist, we measured basal IP accumulation in rat mGlu1a receptor containing CHO-dhfr⁻ cells in the presence of compound A. Figure 1C shows that there is a clear reduction in basal IP production with increasing concentration of compound A. This reduction was statistically significant ($p < 0.05$) as off 1 μ M compound A, at which basal IP accumulation decreased by 24 ± 4 %. A maximal decrease of 33 ± 3 % was found when using 100 μ M compound A. These data indicate that compound A can indeed act as an inverse agonist towards the mGlu1a receptor.

Characterization of [³H]Compound A binding to rat mGlu1a receptor CHO-dhfr⁻ membranes. The specific binding of 2.5 nM [³H]Compound A at 4°C to rat mGlu1a receptor CHO-dhfr⁻ membranes was proportional to the amount of membrane protein and increased linearly between 10 and 50 μ g membrane protein per assay (Figure 2). Non-specific binding was defined using 1 μ M compound 135 as inhibitor. compound 135 was identified as a specific mGlu1 receptor antagonist with a potency of 7.2 ± 1.2 nM ($n=3$) for reversal of glutamate-induced [Ca^{2+}]_i mobilization. Using 20 μ g protein per assay, specific binding of [³H]Compound A was ~92% of the total binding; in typical assay conditions, total and non-specific binding were in the range of 3,800 and 300 DPM, respectively.

Addition of 1.2 mM MgCl₂ and 2 mM CaCl₂ caused a slight increase in specific binding (data not shown). Further addition of NaCl (10-100-300 mM) had no effect. While specific binding decreased by 22% at pH 6, increasing the pH up to 10 had no effect (data not shown).

Association kinetics were measured as described in Materials and Methods.

Decreasing the incubation temperature to 4°C, dramatically enhanced specific binding, whereas a low binding was found at 37°C (Figure 3). The association of [³H]Compound A to membranes was extremely fast. At 4°C, 2 min incubation resulted already in a specific binding corresponding to about 70% of the quantity bound at equilibrium. Maximal binding was reached within 5 min incubation for each incubation temperature. Analysis of the association curves resulted in observed association rate constants (k_{ob}) of 0.6285, 2.571 and 1.523 min⁻¹ at 4°C, 25°C and 37°C, respectively. The kinetics of dissociation were also rapid (Figure 4). At 25°C, the radioligand dissociated within as little as 2 min after 1 μ M compound 135 was added to the reaction tubes. The rapid dissociation kinetics at 25°C did not allow us to calculate an accurate dissociation rate constant (k_{off}). Dissociation occurred more gradual when incubated at 4°C. [³H]Compound A was displaced completely within approximately 45 min after the addition of an excess compound 135. Analysis of the

dissociation curve at 4°C resulted in an k_{off} of 0.1249 min⁻¹. k_{on} ($k_{ob}-k_{off}$ /radioligand concentration) at 4°C was 0.1007 nM⁻¹ min⁻¹.

Ligand saturation experiments were performed at apparent binding equilibrium (30 min incubation) and with 10 concentrations of radioligand. Figure 5 shows the saturation curve and Scatchard Plot of [³H]Compound A binding to rat mGlu1a receptor CHO-dhfr⁻ membranes. Scatchard Plots were linear, indicating the presence of a single, saturable, high affinity binding site. Non-linear regression analysis of the rectangular hyperbola revealed a B_{max} of 6512 ± 1501 fmoles/mg of protein and a K_D of 0.90 ± 0.14 nM (n = 3).

A series of mGlu1 receptor agonists and antagonists was tested for inhibition of [³H]Compound A-binding to rat mGlu1a receptor CHO-dhfr⁻ membranes. Inhibition curves for some antagonists are shown in Figure 6, and the K_i values of all compounds tested are listed in Table 11.

Table 11: Potencies of various mGlu1 receptor agonists and antagonists in inhibition of [³H]Compound A-binding to rat mGlu1a receptor CHO-dhfr⁻ membranes. K_i values and Hill coefficients are mean ± SD of 3-4 independent experiments.

compound	K_i (nM)	Hill coefficient
Compound A	1.35 ± 0.99	0.94 ± 0.04
NPS 2390	1.36 ± 0.50	0.97 ± 0.02
BAY 36-7620	11.2 ± 0.93	0.95 ± 0.02
CPCCOEt	4,900 ± 170	0.93 ± 0.03
glutamate	> 1,000,000	
quisqualate	> 1,000,000	
1S,3R-ACPD	> 1,000,000	
(S)-3,5-DHPG	> 1,000,000	
LY367385	> 1,000,000	
(S)-4C3HPG	> 1,000,000	
AIDA	> 1,000,000	
(S)-4CPG	> 1,000,000	
MCPG	> 1,000,000	

Remarkably, all ligands that bind to the glutamate binding site, i.e. glutamate, quisqualate, 1S,3R-ACPD, (S)-3,5-DHPG, LY-367385, (S)-4C3HPG, (S)-4CPG, MCPG and AIDA did not inhibit [³H]Compound A binding. In contrast, the non-competitive mGlu1 receptor antagonists CPCCOEt, BAY 36-7620, NPS 2390 and compound A inhibited [³H]Compound A binding to rat mGlu1a receptor CHO-dhfr⁻ membranes with potencies, generally consistent with their potencies to inhibit mGlu1a receptor function. compound A and NPS 2390 showed the highest affinity, with a K_i of 1.35 ± 0.99 and 1.36 ± 0.50 nM, respectively. BAY 36-7620 inhibited the binding also at nanomolar concentrations, whereas CPCCOEt displaced at micromolar concentrations.

We also investigated the specificity of [³H]Compound A binding towards the mGlu1 versus mGlu2, -3, -4, -5, and -6 receptors. Using [³H]LY341495, 95, 98 and 40% specific binding was found when using membranes prepared from CHO-dhfr⁻ cells expressing the mGlu2, mGlu3, or mGlu6 receptor, respectively. [³H]MPEP was used as a positive control for the mGlu5 receptor and produced 95% specific binding to rat mGlu5 receptor containing membranes. Total binding of 20 nM [³H]Compound A to membranes prepared from CHO-dhfr⁻ cells expressing the rat mGlu2, -3, -4, -5, or -6 receptor was not higher than the binding to membranes from wild-type CHO-dhfr⁻ cells, nor was it higher than the non-specific binding to rat mGlu1a receptor CHO-dhfr⁻ membranes. Furthermore, specific binding of [³H]Compound A to these membranes was investigated using various blanks: 1 μM compound 135, which is expected to bind to the same site as compound A, glutamate and L-SOP, which bind to the glutamate binding pocket and MPEP that binds to an allosteric site on the mGlu5 receptor (see Table 12). None of these blanks displaced [³H]Compound A. Together, these data demonstrate the specificity of [³H]Compound A for the mGlu1 receptor relative to mGlu2, -3, -4, -5 and -6 receptor subtypes.

Table 12: [³H]Compound A is specific for the mGlu1 receptor relative to the mGlu2, -3, -4, -5 or -6 receptor. The specific binding of 20 nM [³H]Compound A to 40 μg membranes from wild-type (CHO-dhfr⁻) cells or from CHO-dhfr⁻ cells expressing rat mGlu2, -3, -4, -5, or -6 receptors is compared to binding of 10 nM [³H]Compound A to 20 μg rat mGlu1a receptor CHO-dhfr⁻ membranes. Various compounds were used to define non-specific binding. Specific binding (SB) data from rat mGlu1a receptor CHO-dhfr⁻ membranes are the mean ± SD of 3 experiments performed in duplicate.

Other SB data are the mean of duplicate determinations from one experiment (ND= not determined).

SB (fmoles/mg protein)	mGlu1	wild-type	mGlu2	mGlu3	mGlu4	mGlu5	mGlu6
Compound 135 as	6057 ±	65	0	0	82	38	0
blanc	1456	ND	34 ^a	0 ^a	57 ^b	0 ^c	0 ^a
various blancs	ND						

^a 1 mM glutamate was used to determine non-specific binding

5 ^b 0.1 mM L-SOP was used to determine non-specific binding

^c 10 µM MPEP was used to determine non-specific binding

Comparison with [³H]quisqualate binding. Saturation binding experiments were performed using 30 µg protein per incubate and 10 concentrations (1, 2, 5, 10, 20, 40, 10 60, 90, 120 and 150 nM) of the mGlu1 receptor agonist [³H]quisqualate (Figure 7). Fitting of the curves revealed a single binding site with K_D and B_{max} values of 22.0 ± 10 nM and 3912 ± 436 fmoles/mg protein, respectively (n=3). Clearly, [³H]Compound A bound to mGlu1a with a much higher affinity than [³H]quisqualate does. The number of binding sites labelled with [³H]quisqualate was ~60% of the number of binding sites 15 labelled by [³H]Compound A.

The same compounds were evaluated for their inhibitory action on [³H]quisqualate binding to rat mGlu1a receptor CHO-dhfr⁻ membranes. Inhibitory potencies of the tested agonists and antagonists as well as Hill coefficients are summarized in Table 13. 20 In this case, the compounds known to exert a competitive interaction with glutamate, inhibited [³H]quisqualate binding, whereas CPCCOEt, BAY 36-7620 and NPS 2390 did not affect [³H]quisqualate binding. Also compound A did not displace binding of [³H]quisqualate to the rat mGlu1a receptor. The competitive mGlu1 receptor ligands displaced [³H]quisqualate binding with the following rank order of potency: 25 quisqualate > glutamate > LY367385 > (S)-3,5-DHPG > (S)-4C3HPG > 1S,3R-ACPD > (S)-4CPG > AIDA > MCPG.

Table 13: Potencies of various mGlu1 receptor agonists and antagonists in inhibition of [³H]quisqualate binding to rat mGlu1a receptor CHO-dhfr⁻ membranes. K_i values and Hill coefficients are mean ± SD of 2 independent experiments.

compound	K _i (μM)	Hill coefficient
quisqualate	0.030 ± 0.00	0.98 ± 0.01
glutamate	0.40 ± 0.07	0.99 ± 0.01
LY367385	1.18 ± 0.47	0.99 ± 0.01
(S)-3,5-DHPG	1.42 ± 0.00	0.97 ± 0.00
(S)-4C3HPG	1.65 ± 0.06	0.98 ± 0.02
1S,3R-ACPD	1.92 ± 0.20	0.97 ± 0.01
(S)-4CPG	4.51 ± 0.78	0.98 ± 0.00
AIDA	98.3 ± 15.4	0.98 ± 0.03
MCPG	165 ± 0.47	0.96 ± 0.02
Compound A	> 1,000	
NPS 2390	> 1,000	
BAY 36-7620	> 1,000	
CPCCOEt	> 1,000	

5

Nature of competition between CPCCOEt, BAY 36-7620, NPS 2390 and [³H]Compound A binding. The fact that the non-competitive compounds all displaced [³H]Compound A binding without affecting the binding of [³H]quisqualate suggested that these antagonists bound another site than the glutamate binding site. In order to assess whether the reference compounds CPCCOEt, BAY 36-7620, NPS 2390 and the newly identified mGlu1 receptor antagonist compound A compete for the same site or mutually exclusive sites, saturation experiments with [³H]Compound A concentrations from 0.2 to 20 nM in the absence and the presence of CPCCOEt (30 μM), BAY 36-7620 (100 nM) and NPS 2390 (10 nM) were performed. The presence of these competitors did not affect the B_{max} values, but caused a significant increase in the K_D value of [³H]Compound A (Table 14). This is visualized in Figure 8, where the data are plotted using linear regression. In Scatchard plots, the obtained linear lines indeed merge to the same intercept on the X-axis (i.e. the B_{max} value).

10

15

Table 14: K_D and B_{max} values obtained from analyses of [3H]Compound A saturation binding curves obtained in the absence and presence of the mGlu1 receptor antagonists CPCCOEt (30 μM), BAY 36-7620 (100 nM) and NPS 2390 (10 nM). Values are mean \pm SD from 3 individual experiments. Statistical analysis was performed using the Student's t-test (two-tailed): ** $p < 0.01$ and *** $p < 0.001$.

	control	CPCCOEt	BAY 36-7620	NPS 2390
K_D (nM)	0.73 ± 0.09	3.17 ± 0.90 **	5.21 ± 1.2 **	2.90 ± 0.20 ***
B_{max} (fmoles/mg protein)	7284 ± 970	7009 ± 1231	5872 ± 1018	6887 ± 2804

[3H]Compound A binding in rat brain membranes and sections. We used the specific mGlu1 receptor radioligand [3H]Compound A to examine receptor binding in different regions of the rat brain. Membranes from rat cortex, striatum, cerebellum and hippocampus were prepared and [3H]Compound A binding was measured. Non-specific binding compared to total binding was 10% in cerebellum, 30% in hippocampus and 25% in cortex and striatum. K_D and B_{max} values were determined for each brain region (Table 15). K_D values were about 1 nM for all structures. The B_{max} values were significantly different among the various areas. [3H]Compound A labelled a remarkably high number of mGlu1 receptors in the cerebellum. In the striatum and hippocampus about 16% of the number of sites found in the cerebellum were labelled. Only 11% of the number of binding sites in the cerebellum was bound in the rat cortex. Importantly, also incubation with 10 μM of the structurally unrelated compound BAY 36-7620 maximally inhibited [3H]Compound A binding (Figure 9). The mGlu5 receptor selective compound MPEP (tested up to 30 μM) did not affect [3H]Compound A binding to rat cerebellar membranes, again showing the mGlu1 receptor selectivity of compound A.

Using radioligand autoradiography, we examined [3H]Compound A binding distributions in rat brain sections in further detail (Figure 10). [3H]Compound A autoradiography was investigated in sagittal rat brain sections; non-specific binding was determined using compound 135 (Figure 10, panel A). Very high specific binding was observed in the molecular layer of the cerebellum. A moderate signal was observed in the CA3 field and dentate gyrus of the hippocampal formation, thalamus, olfactory tubercle, amygdala and substantia nigra reticulata. The cerebral cortex, caudate putamen, ventral pallidum, nucleus accumbens showed lower labelling. Also

incubation with BAY 36-7620 completely inhibited [³H]Compound A binding to rat brain sections (Figure 10, panel C).

Table 15: Equilibrium binding constants of [³H]Compound A-binding to membranes from rat cortex, hippocampus, striatum and cerebellum. K_D and B_{max} values are mean ± SD derived from 3 independent experiments.

	cortex	hippocampus	striatum	cerebellum
K_D (nM)	1.04 ± 0.40	0.72 ± 0.22	0.84 ± 0.23	0.99 ± 0.36
B_{max} (fmol/mg protein)	471 ± 68	688 ± 125	741 ± 48	4302 ± 2042

Discussion

Up to now, only a few mGlu1 receptor subtype selective antagonists have been found. The mGlu1 receptor has been shown to be selectively blocked by CPCCOEt (Litschig et al., *Mol. Pharmacol.* **55**:453-461, 1999) and BAY 36-7620, with potencies that vary from micromolar for CPCCOEt (6.6 μM) to high nanomolar concentrations for BAY 36-7620 (160 nM). In the present study, compound A is identified as a novel mGlu1 receptor antagonist with low nanomolar functional antagonistic potency on the rat mGlu1a receptor (21.6 nM) and the human mGlu1a receptor (10.4 nM). The antagonist action of compound A was found to be non-competitive, since the maximal glutamate-induced mGlu1 receptor activation was decreased in the presence of compound A. The observed increase in glutamate EC₅₀ in the presence of compound A can be explained by the presence of spare receptors. In the presence of low concentrations of a non-competitive antagonist, the concentrations-response curve will be shifted to the right since more agonist is needed to compensate for the 'nonspare' receptors that are blocked by the antagonist. These antagonist concentrations will not yet affect the maximal agonist response, while higher antagonist concentrations will eventually suppress the maximum response (Zhu et al., *J.Pharm. Tox. Meth.* **29**:85-91, 1993). This phenomenon has also been reported for BAY 36-7620 (Carroll et al., *Mol. Pharmacol.* **59**:965-973, 2001) and CPCCOEt (Hermans et al., *Neuropharmacology* **37**:1645-1647, 1998). Our data further show that compound A may act as an inverse agonist towards the mGlu1a receptor and that compound A acts selectively on the mGlu1 receptor with regard to other mGlu receptor subtypes and ionotropic glutamate receptors. Signal transduction data showed that compound A does not display agonist, antagonist or positive allosteric action on the mGlu2, -3, -4, -5 and -6 receptor and

radioligand binding studies revealed that [³H]Compound A does not bind to the mGlu2, -3, -4, -5 and -6 receptor, furthermore excluding the possibility that compound A acts as a neutral ligand at any of these receptor types. The lack of selective mGlu1 receptor radioligands together with the interesting pharmacological properties of compound A were compelling reasons to label compound A for the investigation of mGlu1 receptors in binding studies.

[³H]Compound A binding met all the requirements for a ligand very well suited to study binding properties, pharmacology and distribution of mGlu1 receptors. First, [³H]Compound A binding studies were performed in rat mGlu1a receptor CHO-dhfr⁻ membranes. Specific binding was very high and increased linearly with protein concentration (Figure 2). Specific binding showed a modest increase in the presence of MgCl₂ and CaCl₂, whereas binding decreased by 22% at pH 6 and was unaffected by an increase in pH. In regard to the effects of pH on binding, it is worth noting the calculated physicochemical properties of compound A: calculated pK_a and clogP are 6.2 and 4.5, respectively. At pH 7.4, the degree of ionisation of compound A is thus very low (only 5.9 %). The percentage of ionisation decreases further at higher pH (1.6 % at pH 8, 0.2 % at pH 9 and no protonation at pH 10). The clogD value remains 4.5 from pH 7.4 to pH 10. At pH 6, however, 61.3 % of compound A is in the protonated form. Accordingly, the clogD decreases to 4.1. The lower binding of the ligand in ionised form suggests that the non-ionised ligand has the highest binding affinity. This is remarkable, and is in contrast with findings for ligands for monoamine G protein-coupled receptors (e.g. the dopamine receptor), which are often strong bases and bind in a cationic form. For such compounds, the driving force for the binding to the receptor is electrostatic in nature (Van de Waterbeemd et al., *J. Med. Chem.* **29**:600-606, 1986). Our data may indicate that ionic interactions are not a driving force in the binding to the receptor, and that there is neither a contribution of ionic surface effects. Additionally, although compound A is a strong lipophilic compound, the very low non-specific [³H]Compound A binding might be due to the fact that no electrostatic interaction can take place between the non-ionised form of compound A and the negatively charged cell membrane. Binding was temperature dependent, and increased substantially at 4°C (Figure 3). By virtue of its fast association and dissociation kinetics, binding equilibrium was rapidly reached. [³H]Compound A labelled apparently a single population of sites with a very high affinity (K_D = 0.90 ± 0.14 nM). In contrast, [³H]quisqualate, the mGlu1 receptor radioligand of choice up to now, exhibited a much higher K_D value of 22.0 ± 10 nM, which correlated well with the value of 37 nM obtained by Mutel et al., *J. Neurochem.*

75:2590-2601, 2000. Besides the considerable higher affinity, [³H]Compound A labelled significantly more (~40%) binding sites than [³H]quisqualate. B_{max} values of 6512 ± 1501 fmoles/mg of protein and 3912 ± 436 fmoles/mg protein were found for [³H]Compound A and [³H]quisqualate, respectively. This discrepancy can be explained on the basis of the G protein coupling of the receptor. Agonists facilitate the coupling of the receptor to the G protein, which results in a receptor conformation with high affinity for agonists. According to this theory, a full agonist such as quisqualate would predominantly label the high affinity or G protein-coupled receptor state. An antagonist would have equal affinity for coupled and uncoupled receptors, and thus for both the high and low affinity states of the receptor. Our finding that the B_{max} for [³H]Compound A is considerably higher than for [³H]quisqualate is in line with this theory.

A striking finding in this study was that the natural agonist glutamate and also quisqualate were unable to inhibit [³H]Compound A binding to rat mGlu1a receptor CHO-dhfr⁻ membranes, whereas CPCCOEt, BAY 36-7620, NPS 2390 and compound A, known as non-competitive antagonists, all inhibited [³H]Compound A binding to the same maximal level (Figure 6). Inhibition of [³H]Compound A binding by the latter compounds followed sigmoidal curves with Hill coefficients of about 1.0 (Table 11), which gave no indication for binding to multiple sites. It is important to mention that although a structurally related analogue was used to define non-specific binding, a similar low non-specific binding was obtained with structurally unrelated compounds such as BAY 36-7620 when used at 1 μM or more in rat mGlu1a receptor CHO-dhfr⁻ membranes (Figure 6). For [³H]quisqualate, all the amino acid-like structures, known as competitive ligands, could displace [³H]quisqualate from its binding site. Inhibitory potencies of quisqualate, glutamate, LY367385, (S)-3,5-DHPG, (S)-4C3HPG, 1S, 3R-ACPD, (S)-4CPG, AIDA and MCPG (Table 13) were in good agreement with the values reported by Mutel et al. *J. Neurochem.* 75:2590-2601, 2000. In contrast, the above non-competitive compounds did not affect its binding. For CPCCOEt, it has been reported that it does not affect [³H]glutamate binding to membranes prepared from rat mGlu1a receptor-expressing cells (Litschig et al., *Mol. Pharmacol.* 55:453-461, 1999). Furthermore, it has been suggested that CPCCOEt does not bind to the glutamate binding site, but interacts with Thr815 and Ala818 in transmembrane domain VII. CPCCOEt is proposed to interfere with receptor signalling by disrupting an intramolecular interaction between the glutamate-bound extracellular domain and the transmembrane domain VII. Carroll et al. in *Mol. Pharmacol.* 59:965-973, 2001 demonstrated that BAY 36-7620 did not displace [³H]quisqualate from the glutamate

binding pocket. Transmembrane helices 4 to 7 were shown to play a crucial role for binding of BAY 36-7620. Our inhibition experiments performed with [³H]Compound A and [³H]quisqualate suggest that CPCCOEt, BAY 36-7620, NPS 2390 bind to the same site as compound A. Saturation experiments using [³H]Compound A in the absence and the presence of 30 μ M CPCCOEt, 100 nM BAY 36-7620 and 10 nM NPS 2390 further support that these compounds bind to the same or mutually exclusive sites. K_D values significantly increased, whereas the B_{max} value was unaltered (Table 14). These results indicate that although the affinity of [³H]Compound A decreases, high concentrations of [³H]Compound A are still able to displace binding of the another compound from its binding site, which is a typical property of a competitive interaction. In conclusion, our data support the notion that CPCCOEt, BAY 36-7620, NPS 2390 and compound A act on a site different from the glutamate binding pocket, presumably they compete for the same transmembrane segment VII.

Previous group I mGlu receptor binding studies in brain were performed using [³H]glutamate or [³H]quisqualate (Schoepp and True, Neurosci. Lett **145**:100-104, 1992; Wright et al., J. Neurochem. **63**:938-945, 1994; Mutel et al., J. Neurochem. **75**:2590-2601, 2000). These radioligands have the disadvantage of labelling more than one type of glutamate receptor. Therefore, selective inhibitors had to be added to the incubation buffer to prevent labelling to other metabotropic or ionotropic glutamate receptor subtypes. To date, there is no radioligand available to specifically study the binding and distribution of the mGlu1 receptor. The specific mGlu1 receptor labelling of [³H]Compound A makes it particularly useful for the investigation of native mGlu1 receptors in rat or human brain. Experiments using rat cortex, hippocampus, striatum and cerebellum membranes revealed that [³H]Compound A specific binding, defined in the presence of 1 μ M compound 135, was high, especially in the cerebellum (only 10 % non-specific binding). Saturation experiments showed that [³H]Compound A again labelled apparently a single binding site with very high affinity. K_D values of about 1 nM were found for all the different brain areas (Table 15). A striking difference in B_{max} values was found: a large population of binding sites was labelled in the cerebellum, whereas in hippocampus, striatum and cortex moderate to low levels of receptor expression were detected.

Because of its specificity, [³H]Compound A proved to be particularly suitable for investigation of mGlu1 receptor distribution in brain sections using radioligand autoradiography. mGlu1 receptor autoradiography revealed that the highest level of mGlu1 specific binding was present in the molecular layer of the cerebellum. The

granule cell layer was very weakly labelled. These results were also found by Mutel et al. in J. Neurochem. **75**:2590-2601, 2000 who investigated group I mGlu receptor distribution using [³H]quisqualate. In the hippocampal formation, the CA3 dendritic field together with the molecular layer of the dentate gyrus showed abundant labelling.

5 The CA1 area showed very weak [³H]Compound A binding, corresponding well with immunohistochemistry data from Lujan et al., Eur.J.Neurosci. **8**:1488-1500, 1996 and Shigemoto et al., J. Neurosci. **17**:7503-7522, 1997, who showed that in CA1 dendritic fields, an antibody specific for the mGlu5 receptor but not a specific mGlu1 receptor antibody yielded intense immunolabelling. Autoradiography experiments using
10 [³H]quisqualate indeed revealed staining in both the CA1 and CA3 region of the hippocampus, indicating binding to both the mGlu1 and mGlu5 receptor, respectively (Mutel et al., J. Neurochem. **75**:2590-2601, 2000). [³H]Compound A binding was also quite high in the thalamus, olfactory tubercle, amygdala and substantia nigra reticulata and was somewhat lower in the cerebral cortex, caudate putamen, nucleus accumbens
15 and ventral pallidum. The same structures were labelled using [³H]quisqualate (Mutel et al., J. Neurochem. **75**:2590-2601, 2000). Also immunocytochemical findings on the cellular localization of the mGlu1a receptor, using an antibody selective for the mGlu1a receptor, were generally consistent with our data (Martin et al., Neuron. **9**:259-270, 1992). Since [³H]Compound A is expected to label all mGlu1 receptor splice variants
20 known to date, the distribution of 1 splice variant may however differ from that of our radiolabel. For example, in the CA3 region and the caudate putamen, which are labelled by the radioligand, mGlu1b receptor but no or little mGlu1a receptor immunoreactivity was found (Martin et al., Neuron. **9**:259-270, 1992; Shigemoto et al., J. Neurosci. **17**:7503-7522, 1997; Ferraguti et al., J. Comp. Neur. **400**:391-407, 1998).
25 An important point in the demonstration of the identity of the [³H]Compound A-labeled sites was the finding that the structurally different compound BAY 36-7620 also fully displaced [³H]Compound A binding to rat brain membranes (Figure 9) as well as to brain sections (Figure 10), providing a good guarantee that the inhibited binding is purely receptor specific and not related to a structural moiety of the radioligand.

30 In this application, we have shown that [³H]Compound A is an excellent radioligand to study mGlu1 receptors in an heterologous expression system, rat brain homogenates and brain sections. We can conclude that because of its minimal non-specific binding, its high binding affinity and marked selectivity, [³H]Compound A is the ligand of
35 choice for further exploration of the mGlu1 receptor. [³H]Compound A opens perspectives for a detailed investigation of subcellular and cellular localization of the

mGlu1 receptor and for the study of the functional role and regulation of the receptor in various areas.

List of Figures

Fig. 1: Antagonist profile of Compound A. Inhibition of glutamate (30 μ M)-induced Ca^{2+} mobilization in CHO-dhfr⁻ cells expressing the rat mGlu1a receptor is shown in Fig. 1A. Data are expressed as percentage of the signal obtained using 30 μ M glutamate, which was set at 100 % and are mean \pm SD of 3 experiments. Fig. 1B shows a concentration-response curve of glutamate alone or together with 20 nM, 30 nM, 60 nM, 100 nM and 1 μ M Compound A. Values are mean \pm SD of triplicate determinations within 1 experiment. An additional experiment showed the same results. Fig. 1C shows basal IP accumulation in the presence of increasing concentrations of Compound A. Values are expressed as percentage of basal IP production in the presence of solvent, which was set as 100 % and are mean \pm SD of 3 experiments performed in quadruplicate.

Fig. 2: Specific [^3H]Compound A binding is linear with amount of membrane protein. 10 to 50 μ g rat mGlu1a receptor CHO-dhfr⁻ membranes was incubated for 30 min on ice with 2.5 nM [^3H]Compound A. Data are expressed as mean \pm SD of triplicate determinations and are from a representative experiment.

Fig. 3: Association time-course curve for [^3H]Compound A binding to rat mGlu1a receptor CHO-dhfr⁻ membranes. Association kinetics were measured by adding 2.5 nM [^3H]Compound A at different times before filtration and was determined at 3 different temperatures. Data are mean \pm SD of 3 independent experiments performed in duplicate.

Fig. 4: Time-course for dissociation of [^3H]Compound A to rat mGlu1a receptor CHO-dhfr⁻ membranes at 4°C and 25°C. Samples were incubated for 30 min at 4°C or 25°C, then an excess of compound 135 was added, followed by rapid filtration at the time indicated for each data point. Values are mean \pm SD of 2 independent experiments performed in duplicate.

Fig. 5: Representative saturation binding curve and Scatchard Plot of [^3H]Compound A binding to rat mGlu1a receptor CHO-dhfr⁻ membranes. Specific binding (SB) was obtained by calculating the difference between total binding (TB) and non-specific binding (BL), measured in the presence of 1 μ M compound 135. For each experiment, data points were determined in duplicate. The experiment was repeated 3 times.

Fig. 6: Inhibition of 2.5 nM [^3H]Compound A binding to rat mGlu1a receptor CHO-dhfr $^-$ membranes by various mGlu receptor antagonists. Data points represent % of total binding and are mean \pm SD of 3-4 individual experiments.

Fig. 7: Representative saturation binding curve and Scatchard Plot of [^3H]quisqualate binding to rat mGlu1a receptor CHO-dhfr $^-$ membranes. For each experiment, data points were determined in duplicate. The experiment was repeated 2 times.

Fig. 8: Saturation binding curves and Scatchard plots of [^3H]Compound A binding to rat mGlu1a receptor CHO-dhfr $^-$ membranes in the absence and presence of CPCCOEt (30 μM), BAY 36-7620 (100 nM) and NPS 2390 (10 nM). The graph shown is a representative of 3 independent experiments. Data are expressed in nM specifically bound. For each experiment, data points were determined in duplicate.

Fig. 9: Inhibition of 2.5 nM [^3H]Compound A binding to rat cerebellar membranes by BAY 36-7620. Data points represent % of total binding and are mean \pm SD of 2 individual experiments.

Fig. 10: [^3H]Compound A binding to sagittal rat brain sections using autoradiography. Panel A is a representative section showing total binding with 1.5 nM [^3H]Compound A. Panel B is a representative and adjacent section showing non-specific binding with 1.5 nM [^3H]Compound A in the presence of 1 μM compound 135. Panel C is a representative section showing non-specific binding with 1.5 nM [^3H]Compound A in the presence of 10 μM BAY 36-7620. Sections from panel A and B were exposed to [^3H]Hyperfilm, while the section from panel C was exposed to a Fuji Imaging plate. Th, thalamus; SNr, substantia nigra reticulata; CA3, CA3 region of the hippocampus; Dg; dentate gyrus of the hippocampus; Cer; cerebellum; Cp, caudate putamen; Cx, cerebral cortex, Ot, olfactory tubercle, Am, amygdala, Vp, ventral pallidum, Na, nucleus accumbens.